

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
13 December 2001 (13.12.2001)

PCT

(10) International Publication Number  
**WO 01/94548 A2**

(51) International Patent Classification<sup>7</sup>: **C12N**

(21) International Application Number: PCT/US01/18326

(22) International Filing Date: 5 June 2001 (05.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/210,286	8 June 2000 (08.06.2000)	US
60/271,758	26 February 2001 (26.02.2001)	US
Not furnished	29 May 2001 (29.05.2001)	US

Nara 630-0101 (JP). **FRANKENBERG, Nicole**; 1224 E. Street #2, Davis, CA 95616 (US). **GAMBETTA, Gregory, A.**; 2516 Temple Drive, Davis, CA 95616 (US). **MONTGOMERY, Beronda, L.**; 728 W. 17th Street, Bloomington, IN 47404 (US).

(74) Agent: **BERLINER, Robert**; Fulbright & Jaworski L.L.P., 29th Floor, 865 South Figueroa Street, Los Angeles, CA 90017-2571 (US).

(81) Designated States (*national*): CA, JP.

(84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).


(71) Applicant: **THE REGENTS OF THE UNIVERSITY OF CALIFORNIA** [US/US]; 1111 Franklin Street, 12th Floor, Oakland, CA 94607-5200 (US).

Published:  
— without international search report and to be republished upon receipt of that report

(72) Inventors: **LAGARIAS, John, Clark**; 1853 Imperial Avenue, Davis, CA 95616 (US). **KOCHI, Takayuki**; 8916-5 Takayama, Daigakusyukusya A-404, Ikoma,

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **HY2 FAMILY OF BILIN REDUCTASES**



(57) Abstract: This invention identifies a novel family of bilin reductases. Designated herein HY bilin reductases, the enzymes of this invention are useful in a wide variety of contexts including but not limited to the conversion of biliverdins to phytobilins and the assembly of holophytochromes or phytofluors.

**WO 01/94548 A2**

5

## HY2 FAMILY OF BILIN REDUCTASES

### CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims benefit of and priority to USSN \_\_/\_\_, filed on May 29, 2001 and entitled HY2 FAMILY OF BILIN REDUCTASES, USSN 60/271,758 filed on February 26, 2001, and to USSN 60/210,286, filed on June 8, 2000, all of which are  
10 incorporated herein by reference in their entirety for all purposes.

### STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

This invention was made, in part, with Government support under Grant Nos: 98-35304-6404 and AMD-9801768 awarded by the United States Department of  
15 Agriculture. The Government of the United States of America may have certain rights in this invention.

### FIELD OF THE INVENTION

This invention relates to the field of phytochromes. In particular, this invention relates to the discovery of a phytochromobilin synthase and a family of related  
20 enzymes from photosynthetic prokaryotes that is are capable of converting a biliverdin into a the phytochrome- and phycobiliprotein chromophore precursors - phytochromobilin, phycocyanobilin and phycoerythrobilin..

### BACKGROUND OF THE INVENTION

The phytochromes comprise a family of biliprotein photoreceptors that  
25 enable plants to adapt to their prevailing light environment (Kendrick and Kronenberg (1994) Kendrick, Pp. 828 in *Photomorphogenesis in Plants*, Dordrecht, The Netherlands: Kluwer Academic Publishers). Phytochromes possess the ability to efficiently photointerconvert between red light absorbing Pr and far red light absorbing Pfr forms, a property conferred by covalent association of a linear tetrapyrrole (bilin or phytobilin) with  
30 a large apoprotein. Phytochromes from cyanobacteria, to green algae and higher plants consist of a well conserved N-terminal polypeptide, roughly 390-600 amino acids in length (see, e.g. U.S. Patent 6,046,014), to which the phytobilin prosthetic group, e.g., phytochromobilin (PΦB) or phycocyanobilin (PCB) is bound.

5                   Phytobilins are linear tetrapyrrole molecules synthesized by plants, algae,  
and cyanobacteria that function as the direct precursors of the chromophores of the light-  
harvesting phycobiliproteins and of the photoreceptor phytochrome (Beale (1993) *Chem.*  
*Rev.* 93: 785–802; Hughes and Lamparter (1999) *Plant Physiol.* 121: 1059–1068). The  
pathways of phytobilin biosynthesis have been elucidated by biochemical fractionation of  
10 plant and algal extracts, by overcoming a blocked step with exogenous putative  
intermediates, and by analysis of linear tetrapyrrole-deficient mutants (Beale and Cornejo  
(1991) *J. Biol. Chem.* 266: 22328–22332; Beale and Cornejo (1991) *J. Biol. Chem.* 266:  
22333–22340; Beale and Cornejo (1991) *J. Biol. Chem.* 266: 22341–22345; Terry *et al.*  
(1993) *Arch. Biochem. Biophys.* 306: 1–15). These studies indicate that the biosynthesis of  
15 phytobilins shares common intermediates with heme and chlorophyll biosynthetic pathways  
to the level of protoporphyrin IX, at which point the latter two pathways diverge by  
metalation with iron or magnesium (Beale (1993) *Chem. Rev.* 93: 785–802). Phytobilins  
are derived from heme, which is converted to biliverdin IXa (BV), the first committed  
intermediate in their biosynthesis. In red algae, cyanobacteria, and plants, this  
20 interconversion is accomplished by ferredoxin-dependent heme oxygenases that are related  
in sequence to the mammalian heme oxygenase (Cornejo *et al.* (1998) *Plant J.* 15: 99–107.;  
Davis *et al.* (1999) *Proc. Natl. Acad. Sci., USA*, 96: 6541–6546; Muramoto *et al.* (1999)  
*Plant Cell* 11: 335–347). Although they catalyze the same reaction, mammalian heme  
oxygenases use an NADPH-dependent cytochrome P450 reductase to generate reducing  
25 power for heme catabolism (Maines (1988) *FASEB J.* 2: 2557–2568).

The metabolic fate of BV differs in mammals, cyanobacteria, and plants,  
with BV being metabolized by different reductases with unique double-bond specificities  
(Figure 1). Mammalian biliverdin IXa reductase (BVR), an NAD(P)H-dependent enzyme  
that catalyzes the two-electron reduction of BV at the C10 methine bridge to produce  
30 bilirubin IXa (BR), was the first of these enzymes to be discovered (Maines and Trakshel  
(1993) *Arch. Biochem. Biophys.* 300: 320–326). A similar enzyme, encoded by the gene  
*bvdR*, was identified in cyanobacteria (Schluchter and Glazer (1997) *J. Biol. Chem.* 272:  
13562–13569). Cyanobacteria and red algae also possess novel ferredoxin-dependent bilin  
reductases for the synthesis of the linear tetrapyrrole precursors of their phycobiliprotein  
35 light-harvesting antennae complexes (Beale and Cornejo (1991) *J. Biol. Chem.* 266: 22328–  
22332; Beale and Cornejo (1991) *J. Biol. Chem.* 266: 22333–22340; Beale and Cornejo

5 (1991) *J. Biol. Chem.* 266: 22341–22345; Cornejo *et al.* (1998) *Plant J.* 15: 99–107).  
Primarily on the basis of studies with the red alga *Cyanidium caldarium*, these investigators  
proposed that the biosynthesis of the two major phycobiliprotein chromophore precursors,  
phycoerythrobilin (PEB) and phycocyanobilin (PCB), utilized two ferredoxin-dependent  
bilin reductases and several double-bond isomerases. The first bilin reductase catalyzes the  
10 two-electron reduction of BV at the C15 methine bridge to produce the BR isomer 15,16-  
dihydrobiliverdin (DHBV), whereas the second bilin reductase catalyzes the conversion of  
15,16-DHBV to 3Z-PEB, a formal two-electron reduction of the C2 and C3 diene system.  
In *C. caldarium*, an additional enzyme mediates the isomerization of 3Z-PEB to 3Z-PCB,  
both of which appear to be isomerized to their corresponding 3E isomers before assembly  
15 with the nascent phycobiliprotein apoproteins (Beale and Cornejo (1991) *J. Biol. Chem.*  
266: 22328–22332; Beale and Cornejo (1991) *J. Biol. Chem.* 266: 22333–22340; Beale and  
Cornejo (1991) *J. Biol. Chem.* 266: 22341–22345).

More recent studies lend support for a similar pathway of PCB and PEB  
synthesis in cyanobacteria (Cornejo and Beale (1997) *Photosynth. Res.* 51: 223–230). In  
20 contrast with mammals and phycobiliprotein-containing organisms, plants and green algae  
reduce BV to 3Z-PFB by the ferredoxin-independent enzyme P $\Phi$ B synthase, which targets  
the 2,3,3<sup>1</sup>,3<sup>2</sup>-diene system for reduction (Terry *et al.* (1995) *J. Biol. Chem.* 270: 11111–  
11118; Woo *et al.* (1997) *J. Biol. Chem.* 272: 25700–25705). In plants, 3Z-PFB is  
isomerized to its 3E isomer, which appears to be the immediate precursor of the  
25 phytochrome chromophore (*Ibid*). The green alga *Mesotaenium caldarium* possesses a  
second bilin reductase activity that catalyzes the reduction of the 18-vinyl group of PFB to  
produce 3Z-PCB (Wu *et al.* (1997) *J. Biol. Chem.* 272: 25700–25705). These investigations  
also revealed that 3E-PCB is the natural phytochrome chromophore precursor in this  
organism.

30 Despite the extensive biochemical analysis of the phytobilin biosynthetic  
pathways in plants, algae, and cyanobacteria, the low levels of bilin reductase expression  
have previously hindered efforts to clone these enzymes.

### SUMMARY OF THE INVENTION

This invention pertains to the isolation and characterization of a family of  
35 bilin reductases (designated herein as the HY2 family). These bilin reductases catalyze the  
conversion of a biliverdin to a phytobilin and can form component (s) of a phytochrome

5 biosynthetic pathway. The bilin reductases of this invention can be used *in vivo* or *in vitro* to simply convert biliverdins to phytobilins or, in conjunction with other enzymes in the phytochrome synthetic pathway to synthesize complete phytochromes and/or phytofluors.

In one embodiment, this invention provides an isolated HY2 family bilin reductase comprising an amino acid consensus sequence as illustrated in Figure 5 or in  
10 Figure 10 and having bilin reductase activity. In certain embodiments, the bilin reductase comprises at least 50% sequence conservation, preferably at least 70% sequence conservation, most preferably at least 90% sequence conservation as shown in Figure 10 and/or at least 80% sequence conservation more preferably at least 100% sequence conservation as shown in Figure 5. In certain preferred embodiments, the bilin reductase is  
15 PebA and/or PebB.

In another embodiment, this invention provides a ferredoxin-dependent bilin reductase comprising at least 15%, preferably at least 20%, more preferably at least 30%, and most preferably at least 50%, at least 75%, at least 90% or at least 95% sequence identity with an enzyme selected from the group consisting of HY2\_ARATH,  
20 YCP2\_SYNPY, YHP2\_PROMA, YHP3\_PROMA, YCP3\_SYNPY, SLR0116, PcyA\_ANASP, PcyA\_NOSPU, PcyA\_SYNY3, PcyA\_SYN8.1, PcyA\_PROME, PebA\_SYNPY, PebA\_SYN8.1, PebA\_PROMA, PebA\_PROME, PebB\_NOSPU, HY2\_ARATH, RCCR\_ARATH, and RCCR\_HORVU, and where, when aligned with HY2, comprises conserved hydrophobic residues at position 137, 157, 158, 256, and 314. In  
25 preferred embodiments, the bilin reductase, when aligned with HY2, comprises a residue selected from the group consisting of Pro-151, Phe-221, Ser222, and ASP-171 and more preferably when aligned with HY2, comprises Pro-151, Phe-221, Ser-222, and ASP-171.

In certain embodiments, the HY2 bilin reductases (HY2 family members) of this invention exclude (proviso out) one or more of the following: hvrcr, atrccr,  
30 rccr\_horvu, rccr\_arath, , ycp2\_synpy, ycp3\_synpy, and HY2.

In still another embodiment, this invention provides an isolated bilin reductase having bilin reductase activity and comprising an amino acid sequence of polypeptide selected from the group consisting of HY2, athy2, slr0116, c362\_anab, ycp2-synpy, ycp3\_synpy, PcyA\_ANASP, PcyA\_NOSPU, PcyA\_SYNY3, PcyA\_SYN81,  
35 PcyA\_PROME, PebA\_SYNPY, PebA\_SYN81, PebA\_PROMA, PebA\_PROME, PebA\_NOSPU, PebB\_SYNPY, PebB\_SYN81, PebB\_PROMA, PebB\_PROME,

- 5   PebB\_NOSPU, HY2\_ARATH, RCCR\_ARATH, and RCCR\_HORVU, or conservative substitutions thereof. In preferred embodiments, the bilin reductase comprises an amino acid sequence of a polypeptide selected from the group consisting of athy2, slr0116, c362\_anab, ycp2-synpy, ycp3\_synpy, PcyA\_ANASP, PcyA\_NOSPU, PcyA\_SYNY3, PcyA\_SYN81, PcyA\_PROMA, PebA\_SYNPY, PebA\_SYN81, PebA\_PROMA,  
10   PebA\_PROMA, PebA\_NOSPU, PebB\_SYNPY, PebB\_SYN81, PebB\_PROMA, PebB\_PROMA, PebB\_NOSPU, HY2\_ARATH, RCCR\_ARATH, and RCCR\_HORVU.

- This invention also provides methods of converting a biliverdin to a phytobilin. The methods involve contacting a bilin reductase of this invention (e.g. an HY2 family bilin reductase) with a biliverdin whereby the biliverdin is converted to a phytobilin.  
15   In certain embodiments, the bilin reductase is a cyanobacterial bilin reductase, and/or an algal bilin reductase, and/or a plant bilin reductase. The bilin reductase can be a recombinantly expressed bilin reductase. The contacting can be *in vivo* or *ex vivo*. In certain embodiments, the contacting is in a cell and the bilin reductase is a heterologous polypeptide. The methods can further comprise contacting the phytychromobilin with a  
20   second bilin reductase to produce a phytyochrome. In certain embodiments, the methods further comprise contacting the phytychromobilin with a second bilin reductase (e.g. PebB) to produce a phytyofluor. In certain embodiments, the bilin reductase is ycp2-snpy and/or ycp3-snpy.

- This invention also provides isolated nucleic acids encoding a bilin reductase  
25   as described herein (e.g. an HY2 family member). Preferred nucleic acids comprise a vector.

- Also are provided cells comprising a heterologous nucleic acid comprising a nucleic acid encoding a bilin reductase (e.g. an HY2 family member) as described herein. Preferred cells include, but are not limited to algal cells, plant cells, yeast cells, bacterial  
30   cells, insect cells, and mammalian cells.

- In still another embodiment, this invention provides a nucleic acid that specifically hybridizes with a nucleic acid encoding any of the bilin reductases described herein under stringent conditions and that encodes a polypeptide having bilin reductase activity. In certain embodiments, the nucleic acids exclude (proviso out) nucleic acids  
35   encoding one or more of the following: hvrcr, atrcr, rccr\_horvu, rccr\_arath, , ycp2\_synpy, ycp3\_synpy, and HY2. Preferred nucleic acids are vectors (e.g. plasmids, cosmids, etc.).

5 In still another embodiment, this invention provides a method of detecting  
expression of a polypeptide. The method comprises providing a cell comprising a nucleic  
acid encoding an apophytochrome; and a nucleic acid encoding a bilin reductase that  
produces a phytybilin that assembles with said apophytochrome to produce a  
holophytochrome or a phytofluor; and detecting an optical signal produced by the  
10 holophytochrome or phytofluor.

This invention also provides a method of producing a photoactive  
holophytochrome. The method involves co-expressing in a cell (*e.g.*, an algal cell, a yeast  
cell, a bacterial cell, a plant cell, an insect cell, a mammalian cell, *etc.*): a heme oxygenase;  
an apophytochrome; and a ferredoxin-dependent bilin reductase; whereby the cell produces  
15 the photoactive holophytochrome and where one or more of the apophytochrome and the  
ferredoxin-dependent bilin reductase are expressed by heterologous nucleic acids. In  
preferred embodiments, the ferredoxin-dependent bilin reductase is an HY2 family bilin  
reductase (*e.g.* HY2, *pcyA*, *etc.*). In a preferred embodiment, the apophytochrome and the  
ferredoxin-dependent bilin reductase are both expressed by heterologous nucleic acids. In  
20 certain embodiments, the heme oxygenase is expressed by a heterologous nucleic acid. In  
certain particularly preferred embodiments, the photoactive holophytochrome is not a  
phytofluor, while in other preferred embodiments, the photoactive holophytochrome is a  
phytofluor. The apophytochrome can be expressed as a fusion protein with a protein that is  
to be labeled with the phytofluor or holophytochrome. In certain preferred embodiments,  
25 the method comprises expressing the ferredoxin-dependent bilin reductase *pebA* and/or  
*pebB*. In a particularly preferred embodiment the cell is a bacterial cell (*E. coli*). The  
method can further involve recovering the photoactive holophytochrome or phytofluor from  
the cell.

In another embodiment this invention provides a cell (*e.g.*, an algal cell, a  
30 yeast cell, a bacterial cell, a plant cell, an insect cell, and a mammalian cell) comprising:  
a heme oxygenase; an apophytochrome; and a ferredoxin-dependent bilin reductase; whereby  
the cell produces a photoactive holophytochrome and where one or more of the  
apophytochrome and the ferredoxin-dependent bilin reductase are expressed by  
heterologous nucleic acids. The ferredoxin-dependent bilin reductase is preferably an HY2  
35 family bilin reductase (*e.g.* HY2, *pcyA*, *etc.*). In certain embodiments, the apophytochrome  
and the ferredoxin-dependent bilin reductase are both expressed by heterologous nucleic

5 acids. They can both be expressed by the same heterologous nucleic acid. In certain cells the heme oxygenase is an endogenous heme oxygenase. In other cells, the heme oxygenase is expressed by a heterologous nucleic acid. The expressed holophytochrome is, in certain  
embodiments, not a phytofluor and, in other embodiments, is a phytofluor. Certain  
preferred cells express *pebA* and/or *pebB*. One preferred cell is a bacterial cell (*e.g.* *E.*  
10 *coli*).

This invention also provides an isolated nucleic acid comprising: a nucleic acid encoding a heme oxidoreductase; and a nucleic acid encoding a ferredoxin-dependent bilin reductase; where the nucleic acid expresses a functional heme  
oxidoreductase and a functional bilin reductase. The heme oxidoreductase and the bilin  
15 reductase can be under control of the same, or different, promoters. The promoter can include a constitutive promoter, an inducible promoter, or a tissue-specific promoter. The nucleic acid can be present in a cell (*e.g.* a bacterial cell, a plant cell, a yeast cell, a mammalian cell, an insect cell, *etc.*). Preferred nucleic acids include one or more genes selected from the group consisting of *HO1*, *HY2*, *PcyA*, *PebA*, and *PebB*. One preferred  
20 nucleic acid comprises an *HO1* coding region and/or a *pcyA* coding region and/or a *pcyB*.

### Definitions.

The term "fluorescent adduct" refers to a fluorescent molecule (*i.e.*, one capable of absorbing light of one wavelength and emitting light of a second wavelength) comprising an "apoprotein" (also referred to as an apophytochrome) component joined to a  
25 "bilin" component, both of which are described below. The fluorescent phytochrome-bilin conjugates (*e.g.*, phytochrome-PEB adducts), are also referred to herein as "phytofluors". The manner in which the two components are joined to form an adduct is irrelevant to the present invention. Typically, the two components spontaneously form an adduct through covalent interactions. The components may also be deliberately linked through covalent  
30 bonds (*e.g.*, through the use of crosslinking reagents). The fluorescent adducts of this invention do not require pairing of an apoprotein with its corresponding native bilin. To the contrary, the invention contemplates adducts consisting of naturally occurring or engineered apoproteins with bilins derived from different organisms, or with non-naturally occurring synthetic linear oligopyrroles or oligopyrrole mimetics.

35 The terms "apoprotein", "apophytochrome", or "apoprotein polypeptide", as used herein, refer to polypeptides derived from eukaryotes, such as vascular plants, non-



5 vascular plants, and algae, or from prokaryotes, such as cyanobacteria and prochlorophytes. The term encompasses both naturally occurring apoproteins and variant polypeptides, *e.g.* derived through mutagenesis. The apoproteins have a hydrophobic pocket, referred to as chromophore binding site, capable of forming an adduct with a bilin component. The apoproteins of the invention are typically homodimeric proteins about 1100 amino acids in  
10 length, each subunit being composed of two major domains. The globular 70 kD N-terminal domain contains the hydrophobic pocket, while the more elongated 55 kD carboxyl terminal domain contains the sites at which the two subunits are associated. Preferred analogues are recognized by and thus comprise the consensus sequence of Figure 6 in U.S. Patent 6,046,014. The apoprotein can be derived from vascular and non-vascular plants,  
15 green alga, or bacteria, can be recombinantly expressed, or can be chemically synthesized *de novo*. Preferred apoproteins are encoded by plant genes, algal genes, bacterial genes, or cyanobacterial genes. Particularly preferred apoproteins include any of the apoproteins described herein or in U.S. Patent 6,046,014 or those listed in the sequence listing of U.S. Patent 6,046,014 or conservative substitutions of these sequences. Most preferred  
20 apoproteins include apoproteins from plants (*e.g.*, oats with an apoprotein having about 1100 amino acid residues), green algae (*e.g.*, *Mesotaenium caldariorum*), or cyanobacteria (as illustrated in U.S. Patent 6,046,014), or related, proteins having conservative substitutions. Truncated apoproteins consisting of a chromophore domain; the apoprotein N-terminal subsequence sufficient for lysase activity are particularly preferred. One  
25 preferred N-terminal subsequence consists of less than about 600 N-terminal amino acids, more preferably less than about 515 N-terminal amino acids, and most preferably less than about 400 N-terminal amino acids. Apophytochromes can be readily identified by one of skill in the art by comparison of the polypeptide sequence in question with the apophytochrome consensus sequence provided in Figure. 6 of U.S. Patent 6,046,014 using  
30 standard sequence comparison methodologies. For a general discussion of apoprotein structure and function, see, Quail *et al.* (1997) *Plant Cell and Environment*, 20: 657-665.

The "bilin" components of the adducts of the invention are linear polypyrroles (*e.g.*, di-, tri-, or tetrapyrroles) capable of fluorescing, or photointerconverting between spectrophotometrically distinct forms, when associated with an apoprotein.  
35 Typically, the bilin components of the invention are isolated from vascular plants, algae, or cyanobacteria according to standard techniques. The bilin components can also be

5 synthesized *de novo*. For a general discussion of bilins useful in the present invention see, Falk (1989) Pp.355-399 In: *The Chemistry of Linear Oligopyrroles and Bile Pigments*, Springer-Verlag, Vienna.

The term "chromophore domain" or "minimal chromophore domain" refers to the apoprotein N-terminal subsequence sufficient for lyase activity; the ability to  
10 spontaneously assemble in the presence of a bilin to form a phytofluor. Chromophore domains typically comprise less than 600 amino acids of the N terminus of the apoprotein, preferably less than about 515 amino acids, more preferably less than about 450 amino acids and most preferably less than about 400, 390, 350 or even as few as 197 N-terminal amino acids known as the "bilin lyase domain", see Wu and Lagarias (2000) *Biochemistry*  
15 39: 13487-13495.. One preferred chromophore domain comprises the 514 N-terminal amino acids of a cyanobacterial phytochrome.

The terms "polypeptide", "peptide" and "protein" are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical analogue of a  
20 corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers. The term also includes variants on the traditional peptide linkage joining the amino acids making up the polypeptide.

The terms "nucleic acid" or "oligonucleotide" or grammatical equivalents herein refer to at least two nucleotides covalently linked together. A nucleic acid of the  
25 present invention is preferably single-stranded or double stranded and will generally contain phosphodiester bonds, although in some cases, as outlined below, nucleic acid analogs are included that may have alternate backbones, comprising, for example, phosphoramidate (Beaucage *et al.* (1993) *Tetrahedron* 49(10):1925) and references therein; Letsinger (1970) *J. Org. Chem.* 35:3800; Sprinzl *et al.* (1977) *Eur. J. Biochem.* 81: 579; Letsinger *et al.*  
30 (1986) *Nucl. Acids Res.* 14: 3487; Sawai *et al.* (1984) *Chem. Lett.* 805, Letsinger *et al.* (1988) *J. Am. Chem. Soc.* 110: 4470; and Pauwels *et al.* (1986) *Chemica Scripta* 26: 1419), phosphorothioate (Mag *et al.* (1991) *Nucleic Acids Res.* 19:1437; and U.S. Patent No. 5,644,048), phosphorodithioate (Briu *et al.* (1989) *J. Am. Chem. Soc.* 111 :2321, O-methylphosphoroamidite linkages (see Eckstein, *Oligonucleotides and Analogues: A Practical Approach*, Oxford University Press), and peptide nucleic acid backbones and  
35 linkages (see Egholm (1992) *J. Am. Chem. Soc.* 114:1895; Meier *et al.* (1992) *Chem. Int.*

- 5 *Ed. Engl.* 31: 1008; Nielsen (1993) *Nature*, 365: 566; Carlsson *et al.* (1996) *Nature* 380: 207). Other analog nucleic acids include those with positive backbones (Denpcy *et al.* (1995) *Proc. Natl. Acad. Sci. USA* 92: 6097; non-ionic backbones (U.S. Patent Nos. 5,386,023, 5,637,684, 5,602,240, 5,216,141 and 4,469,863; Angew. (1991) *Chem. Intl. Ed. English* 30: 423; Letsinger *et al.* (1988) *J. Am. Chem. Soc.* 110:4470; Letsinger *et al.* (1994) *Nucleoside & Nucleotide* 13:1597; Chapters 2 and 3, ASC Symposium Series 580, "Carbohydrate Modifications in Antisense Research", Ed. Y.S. Sanghui and P. Dan Cook; Mesmaeker *et al.* (1994), *Bioorganic & Medicinal Chem. Lett.* 4: 395; Jeffs *et al.* (1994) *J. Biomolecular NMR* 34:17; *Tetrahedron Lett.* 37:743 (1996)) and non-ribose backbones, including those described in U.S. Patent Nos. 5,235,033 and 5,034,506, and Chapters 6 and 7, ASC Symposium Series 580, *Carbohydrate Modifications in Antisense Research*, Ed. Y.S. Sanghui and P. Dan Cook. Nucleic acids containing one or more carbocyclic sugars are also included within the definition of nucleic acids (*see Jenkins et al.* (1995), *Chem. Soc. Rev.* pp169-176). Several nucleic acid analogs are described in Rawls, C & E News June 2, 1997 page 35. These modifications of the ribose-phosphate backbone may be done to facilitate the addition of additional moieties such as labels, or to increase the stability and half-life of such molecules in physiological environments.

The term "heterologous" as it relates to nucleic acid sequences such as coding sequences and control sequences, denotes sequences that are not normally associated with a region of a recombinant construct, and/or are not normally associated with a particular cell. Thus, a "heterologous" region of a nucleic acid construct is an identifiable segment of nucleic acid within or attached to another nucleic acid molecule that is not found in association with the other molecule in nature. For example, a heterologous region of a construct could include a coding sequence flanked by sequences not found in association with the coding sequence in nature. Another example of a heterologous coding sequence is a construct where the coding sequence itself is not found in nature (*e.g.*, synthetic sequences having codons different from the native gene). Similarly, a host cell transformed with a construct which is not normally present in the host cell would be considered heterologous for purposes of this invention.

The terms "isolated" "purified" or "biologically pure" refer to material which is substantially or essentially free from components which normally accompany it as found in its native state. With respect to nucleic acids and/or polypeptides the term can refer to

5 nucleic acids or polypeptides that are no longer flanked by the sequences typically flanking them in nature.

The term "recombinant" or "recombinantly expressed" when used with reference to a cell indicates that the cell replicates or expresses a nucleic acid, or expresses a peptide or protein encoded by a nucleic acid whose origin is exogenous to the cell.

10 Recombinant cells can express genes that are not found within the native (non-recombinant) form of the cell. Recombinant cells can also express genes found in the native form of the cell wherein the genes are re-introduced into the cell by artificial means, for example under the control of a heterologous promoter.

An "HY2-related gene" or a "member of the HY2 family" refers to a gene  
15 that encodes a ferredoxin-dependent bilin reductase and that can catalyze a two or four electron reduction of a linear tetrapyrrole to the biologically active precursors of the chromophores of phytochromes and phycobiliproteins. Typically 200-300 amino acids in length, these enzymes can be recognized by the characteristic signature sequence depicted in Figures 5 and 10.

20 The terms "stringent conditions" or "hybridization under stringent conditions" refers to conditions under which a probe will hybridize preferentially to its target subsequence, and to a lesser extent to, or not at all to, other sequences. "Stringent hybridization" and "stringent hybridization wash conditions" in the context of nucleic acid hybridization experiments such as Southern and northern hybridizations are sequence  
25 dependent, and are different under different environmental parameters. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993) *Laboratory Techniques in Biochemistry and Molecular Biology--Hybridization with Nucleic Acid Probes part I chapter 2 Overview of principles of hybridization and the strategy of nucleic acid probe assays*, Elsevier, New York. Generally, highly stringent hybridization and wash conditions  
30 are selected to be about 5°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Very stringent conditions are selected to be equal to the  $T_m$  for a particular probe.

An example of stringent hybridization conditions for hybridization of  
35 complementary nucleic acids which have more than 100 complementary residues on a filter in a Southern or northern blot is 50% formamide with 1 mg of heparin at 42°C, with the

5 hybridization being carried out overnight. An example of highly stringent wash conditions is 0.15 M NaCl at 72°C for about 15 minutes. An example of stringent wash conditions is a 0.2x SSC wash at 65°C for 15 minutes (*see, Sambrook et al. (1989) Molecular Cloning - A Laboratory Manual (2nd ed.) Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor Press, NY, for a description of SSC buffer*). Often, a high stringency wash is preceded by a 10 low stringency wash to remove background probe signal. An example medium stringency wash for a duplex of, *e.g.*, more than 100 nucleotides, is 1x SSC at 45°C for 15 minutes. An example low stringency wash for a duplex of, *e.g.*, more than 100 nucleotides, is 4-6x SSC at 40°C for 15 minutes. In general, a signal to noise ratio of 2x (or higher) than that observed for an unrelated probe in the particular hybridization assay indicates detection of a 15 specific hybridization. Nucleic acids which do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides which they encode are substantially identical. This occurs, *e.g.*, when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code.

The term conservative substitution is used herein to refer to replacement of 20 amino acids in a protein with different amino acids that do not substantially change the functional properties of the protein. Thus, for example, a polar amino acid might be substituted for a polar amino acid, a non-polar amino acid for a non-polar amino acid, and so forth. The following six groups each contain amino acids that are conservative substitutions for one another: 1) Alanine (A), Serine (S), Threonine (T); 2) Aspartic acid 25 (D), Glutamic acid (E); 3) Asparagine (N), Glutamine (Q); 4) Arginine (R), Lysine (K); 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); and 6) Phenylalanine (F), Tyrosine (Y), and Tryptophan (W).

### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates a pathway for the biosynthesis of bilin pigments. The 30 mammalian bile pigment bilirubin and the linear tetrapyrrole precursors of the phytochrome and phycobiliprotein chromophores of plants, algae, and cyanobacteria share the common intermediate biliverdin IXa. HY2, phytychromobilin synthase or 3Z-phytychromobilin:ferredoxin oxidoreductase; PcyA, 3Z-phytycyanobilin:ferredoxin oxidoreductase; PebA, 15,16-dihydrobiliverdin:ferredoxin oxidoreductase; PebB, 3Z- 35 phytyco-erythrobilin:ferredoxin oxidoreductase; BVR/BvdR, biliverdin IXa:NAD(P)H oxidoreductase. The dashed arrow with a question mark indicates a second type of putative

- 5 3Z-phycoerythrin:ferredoxin oxidoreductase. The dashed arrow indicates a putative phycoerythrin-phycoerythrin isomerase (Beale and Cornejo (1991) *J. Biol. Chem.* 266: 22333-22340).

Figure 2 illustrates phytochrome biosynthesis in *Arabidopsis*.

- Figures 3A and 3B illustrate the *HY2* locus of *Arabidopsis*. Figure 3A shows a map of the region of chromosome 3 containing *HY2*. Two distinct mapping populations were screened, and mapping results with molecular markers are summarized schematically, indicating that *HY2* lies in a region 66 kb in length. Markers starting with the letter c are CAPS markers developed during this study. DNA sequence information for bacterial artificial chromosomes (BACs) MZB10 and F3L24 is available in GenBank/EMBL/DDBJ. The *HY2* gene structure with mutations is illustrated at the bottom. Exons are depicted as dark boxes and thick lines, which reflect coding regions and 59/39 untranslated regions, respectively. Dotted lines indicate introns. Figure 3B shows the genomic sequence of *HY2* and the deduced *HY2* protein sequence from the Columbia (Col) ecotype. Uppercase letters represent exons determined by sequence analysis of *HY2* cDNAs. Introns and spacer sequences are indicated with lowercase letters. The stop codon is double underlined. Mutations in *hy2* alleles are shown in boldface letters. Single nucleotide polymorphisms in both Ler and Wassilewskija (Ws) ecotypes include the following: inserted T (at nucleotide 234), G364T conversion with amino acid change to Asn, and G1182A conversion (silent). Single nucleotide polymorphisms in the Ler ecotype only include the following: C515A (in intron), G884A (silent), C1145T (in intron), and G1717A (in intron). The single nucleotide polymorphism in Ws ecotype only is C1910T (silent).

- Figures 4A and 4B show a RNA gel blot hybridization of MZB10.18/F3L24.1. Figure 4A: Total RNA (10 mg) from 1-week-old seedlings was analyzed by RNA gel blotting using the MZB10.18/F3L24.1 cDNA as a probe. RNA was prepared from seedlings of the *hy2* mutants and corresponding wild-type plants. Figure 4B: The same RNA gel blot was probed with rDNA as a loading control of RNA.

- Figure 5 shows an alignment of *HY2* and *HY2*-Related Proteins. Alignment of the *HY2* protein with proteins of unknown function from oxygenic photosynthetic bacteria identified by PSI BLAST. Conserved residues in 100 or 80% of the aligned sequences are depicted in the consensus sequence with uppercase or lowercase letters,

5 respectively. Sequence similarity groups shown in the consensus sequence reflect conservation in 100% of the sequences. These are labeled as follows: 1, D 5 N; 4, R 5 K; 5, F 5 Y 5 W; and 6, L 5 I 5 V 5 M. Dark shading with white letters, gray shading with white letters, and gray shading with black letters reflect 100, 80, and 60% sequence conservation, respectively. Sequence identifiers correspond to hypothetical proteins from *Synechococcus* 10 sp WH8020 (YCP2\_SYNPY and YCP3\_SYNPY), from *Prochlorococcus* (YHP2\_PROMA and YHP3\_PROMA), and from *Synechocystis* sp PCC 6803 gene (cyanobase locus slr0116; see <http://www.kazusa.or.jp/cyano/cyano.html>). Database accession numbers are AB045112 for HY2 (DDBJ), Q02189 for YCP2\_SYNPY (SWISSPROT), Q02190 for YCP3\_SYNPY (SWISSPROT), CAB95700.1 for YHP2\_PROMA (EMBL), CAB95701.1 15 for YHP3\_PROMA (EMBL), and S76709 for slr0116 (Protein Information Resource). Asterisks are indicated every 20 residues.

Figures 6A through 6D show transient expression of GFP fusion in onion cells and tobacco cells. Figures 6A to 6C: Cells expressing GFP (onion) (Figure 6A), the HY2 chloroplast transit peptide fused to GFP (HY2TP-GFP) (onion) (Figure 6B), and 20 HY2TP-GFP (tobacco) (Figure 6C) were analyzed by fluorescence microscopy using the green channel for GFP. Figure 6D: The same sample as in (C) imaged using the red channel for chlorophyll. Bars in (6A) and (6B) 5 100 mm; bars in (6C) and (6D) 5 10 mm.

Figure 7 shows an SDS-PAGE of the purification of recombinant *Arabidopsis* mHY2. Lane 1, cell-free extract from uninduced *Escherichia coli* strain DH5a carrying 25 pGEXmHY2; lane 2, cell-free extract from isopropylthio- $\beta$ -galactoside-induced *Escherichia coli* strain DH5a carrying pGEXmHY2; lane 3, soluble fraction of the induction; lane 4, GSTmHY2 after glutathione agarose affinity chromatography; lane 5, GSTmHY2 after PreScission protease treatment; lane 6, purified recombinant mHY2 after a second round of glutathioneagarose affinity chromatography; lane M, molecular mass standards. Numbers 30 at right indicate positions of molecular weight markers (Sigma, SDS7) in kilodaltons.

Figure 8 shows phytochrome difference spectra of cph1 after incubation with BV metabolites. A soluble protein extract of isopropylthio- $\beta$ -galactoside2-induced *E. coli* DH5a carrying pGEX-mHY2 was assayed for P $\Phi$ B synthase activity as described herein. Recombinant Cph1 was added to the bilin reaction mixture, which was incubated for 35 another 30 min at room temperature under green safe light, and a phytochrome difference

5 spectrum was obtained. The absorption maximum and minimum are indicated in nanometers.

Figure 9 shows that HY2 converts BV to PΦB as detected by HPLC. Purified recombinant HY2 protein (40 mg) was assayed for PFB synthase activity as described in herein. Bilins were extracted from the incubation mixture using a SepPak C18  
 10 reversed phase cartridge and analyzed by reversed phase HPLC as described herein. The HPLC solvent was acetone:20 mM formic acid (50:50 [v/v]), and the eluate was monitored at 380 nm. The top traces represent the standard bilins BV and 3E-PFB. The third trace shows the bilin metabolites obtained after incubation of BV with HY2. The bottom trace has, in addition, BV as an internal standard.

15 Figure 10 shows a multiple sequence alignment of the identified bilin reductases. All identified sequences were aligned using the programs CLUSTAL W and MEME. Conserved residues in 90 or 70% of the aligned sequences are depicted in the consensus sequence with uppercase or lowercase letters, respectively. Sequence similarity groups, labeled 1 (D, E), 2 (R, K), 3 (F, Y, W), and 4 (L, I, V, M), shown in the consensus  
 20 sequence reflect conservation in >90% of the sequences. Dark shading with white letters, gray shading with white letters, and gray shading with black letters reflect 90, 70, and 50% sequence conservation, respectively. SYNY3, *Synechocystis* sp PCC6803; SYNPY, *Synechococcus* sp WH8020; SYN81, *Synechococcus* sp WH8102; PROMA, *Prochlorococcus* sp SS120; PROME, *Prochlorococcus* sp MED4; NOSPU, *Nostoc punctiforme*; ANASP, *Anabaena* sp PCC7120; ARATH, *Arabidopsis thaliana*; and  
 25 HORVU, *Hordeum vulgare*. Database accession numbers are GB: AF339056 for PcyA\_ANASP (CyanoBase contig 362), GB: AF339057 for PcyA\_NOSPU (JGI contig 632), PIR: S76709 for PcyA\_SYNY3, PcyA\_SYN81 is on JGI contig 51, GB: AF352050 for PcyA\_PROME (JGI contig 26), SW: Q02189 for PebA\_SYNPY, PIR: S31075  
 30 (fragment)/ JGI contig 72 for PebA\_SYN81, EMB: CAB95700.1 for PebA\_PROMA, PebA\_PROME is on JGI contig 26, GB: AF352049 for PebA\_NOSPU (JGI contig 622), SW: Q02190 for PebB\_SYNPY, PebB\_SYN81 is on JGI contig 72, EMB: CAB95701.1 for PebB\_PROMA, PebB\_PROME is on JGI contig 26, GB: AF339058 for PebB\_NOSPU (JGI contig 622), DDBJ: AB045112 for HY2\_ARATH, EMB: CAB77705.1 for  
 35 RCCR\_HORVU, EMB: CAB16763.1 for RCCR\_ARATH. Asterisks indicate every tenth



5 amino acid; dashes indicate gaps; numbers above the line indicate amino acid sequence numbering starting with number one.

Figure 11 shows a phylogenetic tree of the HY2 family of ferredoxin-dependent bilin reductases.

Figure 12 shows an SDS-PAGE of affinity-purified bilin reductases. Lane 1, GST-PcyA\_SYNY3 after glutathione agarose affinity chromatography; lane 2, purified recombinant PcyA\_ANASP after a second round of glutathione agarose affinity chromatography; lane 3, mHY2; lane 4, PebA\_SYNPY; lane 5, PebB\_SYNPY. STD, molecular mass standards (in kilodaltons).

Figures 13A and 13B show phytochrome difference spectra and phytofluor fluorescence spectra of recombinant cyanobacterial phytochrome (Cph1) incubated with reaction metabolites. Figure 13A: BV was incubated with a soluble protein extract of isopropylb thiogalacto pyranoside-induced *E. coli* DH5a strain carrying pGEXNN under standard PFB synthase assay conditions for 30 min at 28°C under green safe light.

Recombinant apoCph1 was added to the reaction and incubated for additional 30 min at room temperature under green safelight, and a phytochrome difference spectrum was obtained. The difference spectrum shown as a solid line was obtained with apoCph1 incubated with PcyA\_SYNY3 metabolites, the spectrum shown in dashed lines was obtained with mHY2 metabolites. Absorption maximum and minimum were indicated as nm. Neither PebA\_SYNPY, PebB\_SYNPY, or a mixture of both was able to form a

25 photoconvertible holophytochrome (no difference spectrum shown). Figure 13B: Phytofluor fluorescence spectra of recombinant cyanobacterial phytochrome (Cph1) incubated with PebA and PebB metabolites. The fluorescence excitation and emission spectra of the phytofluor were obtained after incubation of apoCph1 with the reaction metabolites of PebA\_SYNPY and PebB\_SYNPY. The solid line represents the excitation spectrum monitored with an emission wavelength of 590 nm. The dashed line shows the emission spectrum obtained with an excitation wavelength of 545 nm.

Figure 14 shows HPLC analysis of the BV metabolites of PebA, PebB, PcyA, and HY2 bilin reductases. Forty micrograms of purified protein was incubated at 28°C under green safelight in a total assay volume of 5 mL. The assay system contained an NADPH-regenerating system, spinach ferredoxin-NADP<sup>+</sup>-reductase, spinach ferredoxin, and BSA. The reaction was started by adding 5 mM BV and was stopped by placing the

5 mixture on ice. Bilins were extracted from the incubation mixture using a SepPak C18 reversed-phase column and analyzed by HPLC on a Phenomenex Ultracarb 5mm ODS20 4.6 mm 3 250 mm column with a 4.6 mm 3 30 mm guard column. The HPLC solvent was acetone: 20 mM formic acid (50:50, v/v), and the effluent was monitored at 560 nm for the first 11.5 min and at 380 nm for the remaining time. STDS, mixture of different bilin  
10 standards; HY2, metabolites obtained by mHY2; PcyA, metabolites obtained by PcyA\_SYNY3; PebA, metabolites obtained by PebA\_SYNPY; PebB, metabolites obtained by PebB\_SYNPY; PebA 1 PebB, metabolites obtained by a 1:1 mixture of PebA\_SYNPY and PebB\_SYNPY. Symbols are used for better visualization of peaks. Single symbols indicate the 3E isomer (except 15,16-DHBV and BV) and double symbols indicate the 3Z  
15 isomer, respectively.

Figure 15 shows holophytochrome difference spectrum taken of the protein purified from *E. coli* cells strain LMG194 induced to express apoCph1, HO1 and HY2.

Figure 16 shows a comparison of holophytochrome difference spectra taken of the protein isolated from *E. coli* cells strain LMG194 induced to express apoCph1 with  
20 either HO1 and HY2, or HO1 and PcyA.

### DETAILED DESCRIPTION

This invention pertains to the isolation and characterization of a family of bilin reductases (designated herein as the HY2 family). In certain embodiments, these bilin reductases catalyze the conversion of a biliverdin to a phytobilin and form a component of a  
25 phytochrome biosynthetic pathway. The bilin reductases of this invention can be used *in vivo* or *in vitro* to simply convert biliverdins to phytobilins or, in conjunction with other enzymes in the phytochrome synthetic pathway to synthesize complete phytochromes and/or phytofluors. This invention also pertains to the recombinant synthesis of a phytochrome or phytofluor.

30 The phytochrome chromophore biosynthetic pathway shown in Figure 1 has been elucidated by the classical approach of overcoming a blocked step with exogenous putative intermediates (Terry *et al.* (1993) *Arch. Biochem. Biophys.*, 306:1-15) and by analysis of phytochrome-chromophore deficient mutants (Terry (1997) *Plant Cell Environ.*, 20: 740-745). This pathway shares common intermediates with heme and chlorophyll  
35 biosynthesis to the level of protoporphyrin IX, at which point the latter two pathways diverge by metallation with iron or magnesium (Beale (1993) *Chemical Rev.*, 93:785-802;

5 Porra (1997) *Photochem. Photobiol.*, 65:492-516; Reinbothe and Reinbothe (1996) *Eur. J. Biochem.*, 237:323-343). As shown Figure 1, heme is converted to biliverdin IX $\alpha$  (BV), the first committed intermediate in the biosynthetic pathways of the chromophores of the phytochromes and of the light-harvesting phycobiliproteins in cyanobacteria, red algae and cryptophytes. This reaction is accomplished by a ferredoxin-dependent heme oxygenase in  
10 red algae and cyanobacteria (Rhie and Beale (1995) *Arch. Biochem. Biophys.*, 320:182-194; Cornejo and Beale (1997) *Photosynthesis Res.*, 51:223-230) and by an enzyme in plants that is likely to be similar in structure (Terry (1997) *Plant Cell Environ.*, 20: 740-745). This contrasts with heme oxygenases found in mammalian systems that utilize cytochrome P450  
15 *Pharmacol. and Toxicol.*, 37: 517-554).

As illustrated in Scheme 1 (Fig. 1), the metabolic fate of BV differs in green plants, cyanobacteria and mammals, with BV being metabolized by different reductases with unique double bond specificities. Mammalian biliverdin IX $\alpha$  reductase (BVR), an NAD(P)H-dependent enzyme that catalyzes the reduction at the C10 methine bridge to  
20 produce bilirubin (BR), was the first to be discovered (Singleton and Laster (1965) *J. Biol. Chem.*, 240: 4780-4789). Mammalian BVRs are small soluble enzymes consisting of a single NAD(P)H and bilin binding subunit (Kutty and Maines (1981) *J. Biol. Chem.*, 256: 3956-3962; Maines and Trakshel (1993) *Arch. Biochem. Biophys.*, 300: 320-326). Active recombinant versions of rat and human BVRs have been cloned and expressed in *E. coli*  
25 (Fakhrai and Maines (1992) *J. Biol. Chem.*, 267: 4023-4029; McCoubrey and Maines (1994) *Eur. J. Biochem.*, 222: 597-603; Maines et al. (1996) *Eur. J. Biochem.*, 235: 372-381). The unexpected discovery of the gene *bvdR* in the cyanobacterium *Synechocystis* sp PCC 6803, which encodes a BVR that also catalyzes the NADPH-dependent reduction of the C10 methine bridge of BV (Schluchter and Glazer (1997) *J. Biol. Chem.*, 272: 13562-13569),  
30 has established that this enzyme has ancient evolutionary origins. Interestingly, *bvdR* plays a key role in the regulation of phycobiliprotein biosynthesis in this cyanobacterium since its inactivation leads to reduced accumulation of phycocyanin (Schluchter and Glazer (1997) *J. Biol. Chem.*, 272: 13562-13569).

Cyanobacteria possess additional bilin reductases for the synthesis of the  
35 linear tetrapyrrole precursors of their phycobiliprotein light-harvesting antennae complexes (Cornejo and Beale (1997) *Photosynthesis Research*, 51: 223-230). Based on this

5 investigation and previous studies with the red alga *Cyanidium caldarium* (Beale and  
Cornejo (1991) *J. Biol. Chem.*, 266: 22328-22332; Beale and Cornejo (1991) *J. Biol.*  
*Chem.*, 266: 22333-22340; Beale and Cornejo (1991) *J. Biol. Chem.*, 266: 22341-22345),  
Beale and colleagues have proposed that the biosynthesis of the phycobiliprotein  
chromophore precursors, phycoerythrobilin (PEB) and phycocyanobilin (PCB), involves  
10 two ferredoxin-dependent bilin reductases. The first of these enzymes catalyzes the  
reduction of BV at the C15 methine bridge to give 15,16-dihydrobiliverdin (*i.e.* DHBV  
synthase), while the second reduces 15,16-dihydrobiliverdin (DHBV) at the C2 double bond  
to produce 3Z-PEB (see Scheme 1). In *Cyanidium*, an additional enzyme appears to  
mediate the isomerization of 3Z-PEB to 3Z-PCB, both of which appear to be further  
15 isomerized to their corresponding 3E isomers prior to assembly with the nascent  
phycobiliprotein apoproteins.

By contrast with mammals and phycobiliprotein-containing organisms, BV  
is reduced at the C2 double bond in plants and green algae to yield 3Z-P $\Phi$ B by the  
ferredoxin-dependent enzyme P $\Phi$ B synthase (Terry *et al.* (1995) *J. Biol. Chem.*, 270:11111-  
20 11118; Wu *et al.* (1997) *J. Biol. Chem.*, 272:25700-25705). In both higher and lower plants  
(*e.g.* mosses, ferns), 3Z-P $\Phi$ B and/or its 3E-isomer have been established to be the  
immediate precursor of the phytochrome chromophore (Terry *et al.* (1993) *Arch. Biochem.*  
*Biophys.*, 306: 1-15). Recent studies have established that PCB is the natural phytochrome  
chromophore precursor in the green alga *Mesotaenium caldarium*, and both P $\Phi$ B  
25 synthase and P $\Phi$ B reductase, the enzyme that catalyzes the reductive conversion of 3Z-P $\Phi$ B  
to 3Z-PCB, have been detected in soluble protein extracts from the chloroplast of this  
organism (Wu *et al.* (1997) *J. Biol. Chem.*, 272:25700-25705). While a 3Z to 3E P $\Phi$ B  
isomerases have been hypothesized, this enzyme has not been identified in plant extracts  
(Terry *et al.* (1993) *Arch. Biochem. Biophys.*, 306: 1-15; Beale (1993) *Chemical Rev.*,  
30 93:785-802). The final step of phytochrome chromophore biosynthesis is the covalent  
attachment of P $\Phi$ B or PCB to apophytochrome.

Biochemical analysis of known phytochrome chromophore-deficient  
mutants, which include the *hy1* and *hy2* mutants of *A. thaliana* (Koornneef *et al.* (1980)  
*Zeitschrift fur Pflanzenphysiologie*, 100:147-160; Chory *et al.* (1989) *Plant Cell*, 1:867-880),  
35 the *aurea* and *yg2* mutants of tomato (Koornneef *et al.* (1985) *J. Plant Physiol.*, 120:153-  
165; Van Tuinen *et al.* (1996) *Plant Journal*, 9:173-182; Terry and Kendrick (1996) *J. Biol.*

5 *Chem.*, 271:21681-21686), the *pcd1* and *pcd2* mutants of pea (Weller *et al.* (1996) *Plant Cell*, 8: 55-67; Weller *et al.* (1997) *Plant J.*, 11: 1171-1186), supports the conclusion that these mutations reflect lesions in the structural genes for either heme oxygenase or PΦB synthase (reviewed in Terry (1997) *Plant Cell Environ.*, 20: 740-745). Indeed, the HY1 locus of Arabidopsis has been shown to encode a ferredoxin-dependent heme oxygenase.

10 This invention pertains to the cloning and sequence analysis of HY2 and the demonstration that the HY2 locus encodes phytychromobilin synthase, a ferredoxin-dependent bilin reductase enzyme that converts BV to PΦB. In addition it is demonstrated that protein relatives of HY2 are also biliverdin (BV) reductases.

### **I. HY2 and HY2 family members.**

#### **A) HY2.**

15 The genomic sequence of HY2 and the protein sequence are provided in Figure 3B. Based on cDNA sequence analysis, the *HY2* protein contains 329 residues with a calculated molecular mass of 38.1 kD. At its N terminus, the HY2 protein sequence is rich in serine, with few acidic residues (six serine and one aspartic acid among 45 residues), which suggests a possible transit peptide for localization to plastids (Gravel and von Heijne  
20 (1990) *FEBS Lett.* 261: 455-458). The second amino acid after the initiation methionine is alanine, which is often observed in plastid transit peptides.

The program CHLOROP was also used to predict the transit peptide of HY2, and it indicated that the first 45 amino acid residues of the HY2 protein form a chloroplast transit peptide (Emanuelsson *et al.* (1999) *Protein Sci.* 8: 978-984;  
25 <http://www.cbs.dtu.dk/services/ChloroP/>).

The calculated molecular mass of the mature HY2 protein is 33.0 kD and its predicted pI is 5.66, which are similar to those of PΦB synthase purified from oat seedlings. The HY2 protein has no predicted transmembrane helices, which is also consistent with the  
30 observation that oat PΦB synthase is a soluble protein.

#### **B) The HY2 family and family members.**

Using the HY2 protein sequence as a query sequence, HY2 family members are identified using an iterative PSI-BLAST search of the nonredundant GenBank/EMBL database, *e.g.* using default search parameters (Altschul *et al.* (1997) *Nucleic Acids Res.* 25,

5 3389-3402). No *HY2*-related gene was identified by this search in the nearly complete *Arabidopsis* genome. In contrast, this search identified *HY2*-related sequences from two marine cyanobacteria, *Prochlorococcus marinus* sp. SS120 (EMBL accession numbers CAB95700.1 and CAB95701.1) and *Synechococcus* sp. WH8020 (SWISS-PROT accession numbers Q02189 and Q02190), and a related protein sequence from the cyanobacterium  
10 *Synechocystis* sp. PCC 6803 (cyanobase locus *slr0116*; Protein Information Resource accession number S76709).

Both marine cyanobacteria possess two *HY2*-related ORFs that appear to be part of multigene operons. The *Synechococcus* ORFs, *ycp2\_synpy* and *ycp3\_synpy*, are located within a cluster of genes involved in phycobiliprotein biosynthesis (Wilbanks and  
15 Glazer (1993) *J. Biol. Chem.* 268: 1226-1235), whereas the *Prochlorococcus* ORFs, which we term *yhp2\_proma* and *yhp3\_proma*, are located immediately downstream of a gene related to heme oxygenase (GB:AJ278499.1). These observations suggest that these genes are involved in phycobillin biosynthesis.

Examination of highly conserved residues in the entire *HY2* family and those  
20 within each of the five classes of bilin reductases provides information regarding residues important to the protein structure, ferredoxin interaction site, and substrate/product specificity. In this regard, only a small number of residues are conserved in the entire *HY2* family of enzymes. These include hydrophobic residues at positions 137, 157, 158, 256, and 314, Pro151, Phe221, Ser222, and Asp171 (Figure 10). The notable lack of conserved  
25 basic residues suggests that the propionyl moieties of the bilin substrates do not form salt linkages with the enzymes. The conserved hydrophobic residues proline and phenylalanine are likely to be involved in overall protein structure (*i.e.*, folding). Alternately, they may form hydrophobic interactions with conserved regions of the various bilin substrates.

The loss-of-function *hy2-1* and *hy2-104* alleles of phytychromobilin synthase  
30 from *Arabidopsis* support the critical role of Pro151 in *HY2*'s structure. The conserved serine and aspartate residues likely play catalytic roles, such as hydrogen bonding with the substrate and/or substrate protonation to make the bound bilin a better electron acceptor. Despite the wide divergence of the *HY2* family, we believe that these conserved residues indicate that the active sites of all members of this class are similar. We believe the distinct  
35 doublebond reduction specificities of the BV reductases (*i.e.*, PcyA, PebA, *HY2*), the 15,16-DHBV reductases (*i.e.*, PebB), and the RCCR families reflect the positioning of the

5        respective substrates within the catalytic pocket. Because the A/B and C/D rings of BV are  
very similar but not identical, it is conceivable that the substrate binding sites of the PebA  
and HY2 enzymes are tailored to position BV in opposite orientations, favoring electron  
transfer to the bilin C/D ring or A ring, respectively. If this is true, then the PebB class  
might tether its 15,16-DHBV substrate in an orientation similar to that of the HY2 family,  
10        whereas RCC might be bound to RCCR in a manner similar to that in which BV is bound to  
PebA.

### 1)        pcyA

We have documented that the *pcyA* genes of the cyanobacteria *Synechocystis*  
sp PCC6803, *Anabaena* sp PCC7120, and *Nostoc punctiforme* encode bilin reductases that  
15        catalyze the four electron reduction of BV to 3Z-PCB. PCB is the precursor of the  
chromophores of the phycobiliproteins phycocyanin and allophycocyanin, which are  
abundant in all three cyanobacteria. PcyA enzymes are atypical bilin reductases because all  
others catalyze two-electron reductions. Formally, these enzymes catalyze two electron  
reductions of both the A and D rings of BV; however, we have not detected the production  
20        of semireduced intermediates such as PΦB and 18<sup>1</sup>,18<sup>2</sup>-DHBV. Thus, it appears that the  
partially reduced intermediates are tightly bound to the enzyme. The direct conversion of  
BV to PCB in these cyanobacteria is in contrast to the proposed pathways of PCB  
biosynthesis in the red alga *C. caldarium*, which involves the intermediacy of PEB, and in  
the green alga *M. caldariorum*, in which 3Z-PΦB is an isolable intermediate. *pcyA*-related  
25        genes also are present in the oxyphotobacterium *Prochlorococcus* sp. MED4, an  
unanticipated observation in view of the lack of phycobiliproteins in this organism. We  
were able to clone the *Prochlorococcus* sp. MED4 *pcyA* gene and express it as an N-  
terminal GST fusion. We determined that recombinant PcyA\_PROME was able to reduce  
BV to PCB in our standard phytochrome-based assay (data not shown). It therefore  
30        possesses the same enzymatic activity as other studied PcyA enzymes.

### 2)        pebA and pebB

We have observed that the *pebA* and *pebB* genes of the cyanobacteria  
*Synechococcus* sp WH8020 and *N. punctiforme* encode bilin reductases that catalyze the  
conversions of BV to 15,16-DHBV and 15,16-DHBV to 3ZPEB, respectively (Figure 1).  
35        PebA therefore is a 15,16-DHBV:ferredoxin oxidoreductase, whereas PebB is a 3Z-

5 PEB:ferredoxin oxidoreductase. Both activities are consistent with the pathway of PEB biosynthesis in the red alga *C. caldarium*. The two *peb* genes also are found in the same operon in both phycoerythrin-producing cyanobacteria, and their close association with the major phycobiliprotein gene clusters supports their role in phycobilin biosynthesis.

Without being bound to a particular theory, we believe PebA and PebB  
10 function as a dual enzyme complex, in view of the synergistic metabolism of BV observed when the two enzymes are coincubated. A *peb* operon is not present in the genome of the cyanobacterium *Synechocystis* sp PCC6803, an organism that lacks phycoerythrin. This strongly suggests that PCB is synthesized in this cyanobacterium via the PcyA-dependent pathway, as opposed to the PEB pathway found in *C. caldarium*. In this regard,  
15 biochemical analyses of crude extracts from *Synechocystis* sp PCC6803 provide no evidence for the production of PEB. The MED4 and SS120 subspecies of the oxyphotobacteria *Prochlorococcus* also possess *peb* operons very similar to those of *Synechococcus* sp WH8020 and WH8102, except that the former possess upstream genes related to heme oxygenase. This strongly suggests that both oxyphotobacterial subspecies  
20 can synthesize PEB.

We also believe that *Prochlorococcus* PebA and PebB are likely functional orthologs of the *Synechococcus* and *Nostoc* enzymes. It is likely that numerous bilin isomerases are present in these oxygenevolving photosynthetic organisms.

### **C) Identification of other members of the HY2 family.**

25 Other members of the HY2 family of bilin reductases can readily be identified using the methods described herein (*see e.g.*, Example 2). In a preferred embodiment, such methods involve using alignment algorithms with one or more members of the HY2 family as described herein to search nucleic acid and/or protein databanks to identify related genes/polypeptides.

30 The activity of the putative bilin reductase can be confirmed, *e.g.* using a standard bilin reductase activity assay. One such bilin reductase assay is described in detail in Examples 1 and 2. Basically, the putative bilin reductase is combined with a biliverdin in a buffer system compatible with enzyme activity. The assay mixture is incubated for a period of time. Product analysis can be accomplished using a direct HPLC assay (*see*



5 Example 1) or by a coupled assay after the addition of an appropriate apophytochrome (*e.g.* recombinant cyanobacterial phytochrome such as Cph1) using spectroscopic methods.

## **II. Uses for HY2.**

The HY2 bilin reductases of this invention are useful tools for applications. The ability to engineer the biosynthesis of phycoerythrobilin (PEB) in any biliverdin-  
10 producing organism is now feasible via the introduction of one or two genes. Similarly, photoactive holophytochromes (*e.g.* bilin pigments bound to apophytochromes) can be produced in any ferredoxin-containing organism.

Coexpression of bilin reductase genes with apophytochromes enables us to produce holophytochromes in a wide number of cell types including, but not limited to algal  
15 cells, plant cells, bacteria, yeast, vertebrate cells (including mammalian cells), insect cells, and the like. This facilitates not only three-dimensional structural analysis of phytochrome, but also the reconstruction of phytochrome signaling in a non-plant system in which we can exploit the power of molecular genetic analyses. Recombinantly expressed phytochromes thus present an excellent model system useful for a wide variety studies. Similar  
20 approaches has proven invaluable for the structure–function analysis of the steroid hormone receptor family.

By introducing the *pcyA* gene into wild-type and chromophore-deficient mutant plants, it is possible to change the wavelength specificity of phytochrome, which can favorably alter plant growth and development in the field environment. Introduction of the  
25 *pebA* and *pebB* genes into plants can shunt the conversion of BV to PEB, yielding photomorpha-genetically challenged plants with fluorescent phytochromes. This is especially useful for the analysis of the temporal and spatial patterns of phytochrome expression in plants.

### **A) In vivo and ex vivo conversion of biliverdin to phytobilin.**

30 In certain embodiments, the HY2 family of bilin reductases of this invention can be used as simple reagents (reducing agents) to convert a biliverdin to a phytobilin. The enzymes can be used *in vivo* (*e.g.* in a plant) *in vitro* (*e.g.* in a cell culture), or *ex vivo* as a simple reagent. Thus, for example, one or more bilin reductases can be contacted with a biliverdin *ex vivo* in an appropriate buffer system (*e.g.*, typically in the presence of a  
35 ferredoxin) resulting in the conversion of the biliverdin to a phytobilin (*see, e.g.*, Example

- 5 1). The phyto bilin is then readily purified *e.g.* using HPLC, *e.g.* as described herein in the Examples.

Alternatively, a host cell can be transfected with a nucleic acid encoding one or more bilin reductases of this invention and/or other components of the phytochrome biosynthetic pathway. The bilin reductases are expressed in the host cell where, in the presence of ferredoxin, they convert a biliverdin to a phyto bilin. Such methods can be simply used to produce a phyto bilin, or can be used in increase expression/production of a holophytochrome (*e.g.* by augmenting the phytochrome synthetic machinery already present in a plant cell, algal cell, photoactive bacterial cell, *etc.*) or a phytofluor.

Preferred host cells are cells that natively provide a heme and/or a heme oxygenase and/or a ferredoxin. Various preferred cells include, in certain embodiments, cells that do not normally produce a phytochrome (*e.g.* certain bacterial cells, mammalian cells, *etc.*) and in certain other embodiments, cells that typically express phytochromes (*e.g.* plant cells, algal cells, *etc.*).

#### **B) Expression of holophytochromes.**

20 The bilin reductases, and other enzymes identified herein, can be used to assemble photoactive holophytochromes including photoactive chromophore precursors and fluorescent phytofluor chromophore precursors. It was a surprising discovery of this invention that a cell transfected with nucleic acids encoding the components of a bilin synthetic pathway (*e.g.*, HO1, PcyA, and/or HY2) and a nucleic acid encoding an apophytochrome (*e.g.* Cph1, native and recombinant oat phytochrome A (ASPHYAST), *Avena sativa* phyA (AsphyA), *Arabidopsis* phyA (AtphyA), *Mesotaenium caldariorum* phylb (McpHYb), *Synechocystis* sp 6803 phy1 (S6803 phy1), and the like (*see*, U.S. Patent 6,046,014)) will express a phytochromobilin that assembles with the apophytochrome to produce a photoactive holophytochrome (*e.g.* chromophore or phytofluor).

30 The holophytochrome, whether chromophore or phytofluor finds a number of uses. In one particularly preferred use, the chromophore or phytofluor are useful as detectable labels (*e.g.* colorimetric or fluorescent labels). Such labels are useful for the visualization, and/or localization and/or isolation of attached ligands. In particularly preferred embodiments, the apophytochrome is expressed as a fusion protein with a polypeptide that it is desired to label. The apophytochrome can be directly fused to the

35

5 polypeptide or separated by a peptide linker. When the fusion protein is expressed, the apophytochrome component combines with the bilin to produce a chromophore or phytofluor which then acts as a label for the polypeptide.

In particularly preferred embodiments, the holophytochrome is a phytofluor. Phytofluors are fluorescent apophytochrome-bilin conjugates (*e.g.*, apophytochrome-PEB adducts), that are intensely fluorescent, photostable proteins useful as fluorescent labels (10 *e.g.* as probes for biological research, *see, e.g.*, U.S. Patent 6,046,014).

In certain embodiment the host cells are transfected with the *pebA* and *pebB* genes to shunt the conversion of biliverdin to PEB, yielding a fluorescent phytofluor.

### **C) Heterologous holophytochromes as model systems.**

15 The methods of this invention can be used to express a holophytochrome in essentially any cell including cells that, in their native state, do not harbor a phytochrome. Cells containing recombinant holophytochrome provide model systems having a wide variety of uses. For example, such cells can be used to screen for agents that alter the activity and/or spectral sensitivity of the phytochrome. In such assays the cells are (20 contacted with the agent(s) in question and then assayed for changes in physiological activity and/or changes in phytochrome localization or conformation and/or changes in spectral characteristics.

Such model systems are also useful for dissecting the metabolic pathways in which phytochromes are involved.

25 Recombinant holophytochromes of this invention can be introduced into plants, algae, and the like that normally harbor phytochromes as well. Such introduced heterologous phytochromes alter the wavelength specificity plant, which can favorably alter plant growth and development in the field environment. Using such methods the host range of various plants can be improved.

### **III. Cloning and expression of HY2 proteins and other enzymes in the phytochrome biosynthetic pathway..**

It is often desirable to provide isolated ferredoxin-dependent bilin reductases (*e.g.* HY2 family members) and/or holophytochromes (*e.g.* chromophores or phytofluors) of this invention. These polypeptides and/or phytochromes can be used to raise an immune (35 response and thereby generate antibodies specific to the phytochrome or to components of

5 the biosynthetic system which can then be used to localize and/or visualize such elements in cells. In addition, as indicated above, the isolated phytochromes can be coupled to various moieties and act as detectable labels. The enzyme components of the bilin synthetic pathway can be used as chemical reagents.

As explained below, the holophytochromes and components of the  
10 phytochrome and/or bilin synthetic pathway can be conveniently produced using synthetic chemical syntheses or recombinant expression methodologies. In addition to the intact full-length polypeptides, in some embodiments, it is often desirably to express immunogenically relevant fragments (e.g. fragments that can be used to raise specific antibodies).

**A) De novo chemical synthesis.**

15 The phytochrome pathway components and/or apophytochromes the active bilin lyase domain or other subsequences can be synthesized using standard chemical peptide synthesis techniques. Where the desired subsequences are relatively short (e.g., when a particular antigenic determinant is desired) the molecule may be synthesized as a single contiguous polypeptide. Where larger molecules are desired, subsequences can be  
20 synthesized separately (in one or more units) and then fused by condensation of the amino terminus of one molecule with the carboxyl terminus of the other molecule thereby forming a peptide bond.

Solid phase synthesis in which the C-terminal amino acid of the sequence is attached to an insoluble support followed by sequential addition of the remaining amino  
25 acids in the sequence is the preferred method for the chemical synthesis of the polypeptides of this invention. Techniques for solid phase synthesis are described by Barany and Merrifield, *Solid-Phase Peptide Synthesis*; pp. 3-284 in *The Peptides: Analysis, Synthesis, Biology. Vol. 2: Special Methods in Peptide Synthesis, Part A.*, Merrifield, et al. (1963) *J. Am. Chem. Soc.*, 85: 2149-2156, and Stewart et al. (1984) *Solid Phase Peptide Synthesis*,  
30 2nd ed. Pierce Chem. Co., Rockford, Ill.

## **B) Recombinant expression.**

In a preferred embodiment, the holophytochromes of this invention and/or components of the phytochrome synthetic pathway (e.g. HY2 family reductases) are synthesized using recombinant expression systems. Generally this involves creating a DNA sequence that encodes the desired protein(s), placing the DNA in an expression cassette under the control of a particular promoter, expressing the protein in a host, and, if desired isolating the expressed protein.

### **1) Nucleic acids.**

Using the information provided herein, (e.g. HY2 family member sequences, primers, etc.) the nucleic acids (e.g., encoding apoproteins, HY2 family reductases, and the like) can be prepared using standard methods known to those of skill in the art. For example, the HY2 family nucleic acid(s) may be cloned, or amplified by *in vitro* methods, such as the polymerase chain reaction (PCR), the ligase chain reaction (LCR), the transcription-based amplification system (TAS), the self-sustained sequence replication system (SSR), etc. A wide variety of cloning and *in vitro* amplification methodologies are well-known to persons of skill. Examples of techniques sufficient to direct persons of skill through *in vitro* amplification methods are found in Berger, Sambrook, and Ausubel, as well as Mullis *et al.*, (1987) U.S. Patent No. 4,683,202; *PCR Protocols A Guide to Methods and Applications* (Innis *et al.* eds) Academic Press Inc. San Diego, CA (1990) (Innis); Arnheim & Levinson (October 1, 1990) *C&EN* 36-47; *The Journal Of NIH Research* (1991) 3: 81-94; (Kwoh *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86: 1173; Guatelli *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87, 1874; Lomell *et al.* (1989) *J. Clin. Chem.*, 35: 1826; Landegren *et al.*, (1988) *Science*, 241: 1077-1080; Van Brunt (1990) *Biotechnology*, 8: 291-294; Wu and Wallace, (1989) *Gene*, 4: 560; and Barringer *et al.* (1990) *Gene*, 89: 117.

DNA encoding desired proteins (e.g. HY2 family members) described herein can be prepared by any suitable method as described above, including, for example, cloning and restriction of appropriate sequences or direct chemical synthesis by methods such as the phosphotriester method of Narang *et al.* (1979) *Meth. Enzymol.* 68: 90-99; the phosphodiester method of Brown *et al.* (1979) *Meth. Enzymol.* 68: 109-151; the diethylphosphoramidite method of Beaucage *et al.* (1981) *Tetra. Lett.*, 22: 1859-1862; and the solid support method of U.S. Patent No. 4,458,066.

5                   Chemical synthesis produces a single stranded oligonucleotide. This may be converted into double stranded DNA by hybridization with a complementary sequence, or by polymerization with a DNA polymerase using the single strand as a template. One of skill would recognize that while chemical synthesis of DNA is limited to sequences of about 100 bases, longer sequences may be obtained by the ligation of shorter sequences.

10                   Alternatively, subsequences may be cloned and the appropriate subsequences cleaved using appropriate restriction enzymes. The fragments may then be ligated to produce the desired DNA sequence.

                  In one embodiment, the nucleic acids of this invention can be cloned using DNA amplification methods such as polymerase chain reaction (PCR) (*see, e.g., Example*  
15 1). Thus, for example, the nucleic acid sequence or subsequence is PCR amplified, using a sense primer containing one restriction site (*e.g., NdeI*) and an antisense primer containing another restriction site (*e.g., HindIII*). This will produce a nucleic acid encoding the desired sequence (*e.g. HY2 sequence*) or subsequence and having terminal restriction sites. This nucleic acid can then be easily ligated into a vector containing a nucleic acid encoding the  
20 second molecule and having the appropriate corresponding restriction sites. Suitable PCR primers can be determined by one of skill in the art using the sequence information and representative primers are provided herein. Appropriate restriction sites can also be added to the nucleic acid encoding the desired protein or protein subsequence by site-directed mutagenesis. The plasmid containing the desired sequence or subsequence (*e.g. HY2 bilin*  
25 reductase sequence) is cleaved with the appropriate restriction endonuclease and then ligated into the vector encoding the second molecule according to standard methods.

                  The nucleic acid sequences encoding desired protein or protein subsequences may be expressed in a variety of host cells, including *E. coli*, other bacterial hosts, yeast, and various higher eukaryotic cells such as the COS, CHO and HeLa cells lines and  
30 myeloma cell lines. The recombinant protein gene will be operably linked to appropriate expression control sequences for each host. For *E. coli* this includes a promoter such as the T7, trp, or lambda promoters, a ribosome binding site and preferably a transcription termination signal. For eukaryotic cells, the control sequences will include a promoter and often an enhancer (*e.g., an enhancer derived from immunoglobulin genes, SV40,*  
35 cytomegalovirus, *etc.*), and a polyadenylation sequence, and may include splice donor and acceptor sequences.

5                   The isolation and expression of an HY2 nucleic acid is illustrated in  
Examples 1 and 2.

## 2) Expression Vectors

A nucleic acid of the invention encoding a one or more enzymes of a  
phytochrome biosynthetic pathway, *e.g.*, as described above, can be incorporated into a  
10 recombinant expression vector in a form suitable for expression of the enzyme(s) (and in  
certain embodiments, assembly of a holophytochrome) in a host cell. The term "in a form  
suitable for expression of the fusion protein in a host cell" is intended to mean that the  
recombinant expression vector includes one or more regulatory sequences operably linked  
to the nucleic acid encoding the enzyme(s) in a manner that allows for transcription of the  
15 nucleic acid into mRNA and translation of the mRNA into the subject protein(s). The term  
"regulatory sequence" is art-recognized and intended to include promoters, and/or enhancers  
and/o other expression control elements (*e.g.*, polyadenylation signals). Such regulatory  
sequences are known to those skilled in the art (*see, e.g.*, Goeddel (1990) *Gene Expression  
Technology: Meth. in Enzymol.* 185, Academic Press, San Diego, CA; Berger and Kimmel,  
20 *Guide to Molecular Cloning Techniques, Methods in Enzymology* 152 Academic Press, Inc.,  
San Diego, CA; Sambrook *et al.* (1989) *Molecular Cloning - A Laboratory Manual* (2nd  
*ed.*) Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor Press, NY,,*etc.*).

The design of the expression vector may depend on such factors as the  
choice of the host cell to be transfected and/or particular protein(s) to be expressed. When  
25 used in mammalian cells, a recombinant expression vector's control functions are often  
provided by viral genetic material. Preferred promoters include, but are not limited to CMV  
immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and  
mouse metallothionein-I. Use of appropriate regulatory elements can allow for high level  
expression of the polypeptide(s) in a variety of host cells. A number of suitable expression  
30 systems are commercially available. In one preferred embodiment, the sequences encoding  
the desired polypeptide(s) are expressed in TA cloning plasmid, pCR2.1 (Invitrogen), *e.g.* as  
described in Example 3.

It will be appreciated that desired polypeptides can be operably linked to  
constitutive promoters for high level, continuous expression. Alternatively, inducible  
35 and/or tissue-specific promoters can be utilized.

5 In one embodiment, the recombinant expression vector of the invention is a plasmid or cosmid. Alternatively, a recombinant expression vector of the invention can be a virus, or portion thereof, which allows for expression of a nucleic acid introduced into the viral nucleic acid. For example, replication defective retroviruses, adenoviruses and adeno-associated viruses can be used.

10 Examples of techniques and instructions sufficient to direct persons of skill through cloning procedures are found in Berger and Kimmel, *Guide to Molecular Cloning Techniques, Methods in Enzymology* 152 Academic Press, Inc., San Diego, CA; Sambrook *et al.* (1989) *Molecular Cloning - A Laboratory Manual (2nd ed.) Vol. 1-3*, Cold Spring Harbor Laboratory, Cold Spring Harbor Press, NY; Ausubel *et al.* (1994) *Current Protocols in Molecular Biology*, Current Protocols, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., U.S. patent number 5,017,478; and European Patent No. 0,246,864.

### 3) Host cells.

20 The holophytochromes and/or components of the phytochrome biosynthetic pathway can be expressed in virtually any cell. Preferred cells, however, comprise an endogenous heme and/or a ferredoxin or are modified to comprise a heme and/or a ferredoxin. Particularly preferred cells include, but are not limited to algal cells, bacterial cells, yeast cells, plant cells, vertebrate cells, and mammalian cells including human cells.

25 The holophytochromes and/or components of the phytochrome biosynthetic pathway are expressed in a host cell by introducing nucleic acid encoding the subject polypeptide(s) into the host cell, wherein the nucleic acid is in a form suitable for expression of the subject polypeptide(s) in the host cell. For example, a recombinant expression vector of the invention, encoding the subject polypeptide(s), is introduced into a host cell. Alternatively, nucleic acid encoding the subject polypeptide(s) which is  
30 operatively linked to regulatory sequences (*e.g.*, promoter sequences) but without additional vector sequences can be introduced into a host cell.

As used herein, the term "host cell" is intended to include any cell or cell line so long as the cell or cell line is not incompatible with the protein(s) to be expressed, the selection system chosen or the culture system employed. As indicated above suitable cells  
35 include, but are not limited to algal cells, bacterial cells (*e.g.* *E. coli*), yeast cells (*e.g.*, *S. cerevisiae*, *S. pombe*, *P. pastoris*, *K. lactis*, *H. polymorpha*, *see, e.g.*, Fleer (1992) *Curr.*



5 *Opin. Biotech.* 3(5): 486-496), fungal cells, plant cells (*e.g. Arabidopsis*), invertebrate cells  
(*e.g. insect cells*) and vertebrate cells including mammalian cells. Non-limiting examples of  
mammalian cell lines which can be used include CHO cells (Urlaub and Chasin (1980)  
*Proc. Natl. Acad. Sci. USA* 77: 4216-4220), 293 cells (Graham *et al.* (1977) *J. Gen. Virol.*  
36: 59), or myeloma cells like (*e.g.*, SP2 or NS0, *see* Galfre and Milstein (1981) *Meth.*  
10 *Enzymol.* 73(B):3-46), and the like.

Examples of vectors for expression in yeast *S. cerevisiae* include, but are not  
limited to pYepSec1 (Baldari. *et al.* (1987) *Embo J.* 6: 229-234), pMFa (Kurjan and  
Herskowitz, (1982) *Cell* 30: 933-943), pJRY88 (Schultz *et al.*, (1987) *Gene* 54:113-123),  
and pYES2 (Invitrogen Corporation, San Diego, Calif.). The desired polypeptides can be  
15 expressed in insect cells (*e.g.*, SF9 cells) using baculovirus expression vectors (*see, e.g.*,  
O'Reilly *et al.* (1992) *Baculovirus Expression Vectors: A Laboratory Manual*, Stockton  
Press).

#### **4) Introduction of nucleic acid into a host cell.**

Nucleic acid(s) encoding the apophytochrome and/or components of the bilin  
20 biosynthetic pathway n can be introduced into a host cell by standard techniques for  
transfecting cells. The term "transfecting" or "transfection" is intended to encompass all  
conventional techniques for introducing nucleic acid into host cells, including calcium  
phosphate co-precipitation, DEAE-dextran-mediated transfection, lipofection,  
electroporation and microinjection. Suitable methods for transfecting host cells can be  
25 found in Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual, 2nd Edition*,  
Cold Spring Harbor Laboratory press, and other laboratory textbooks.

The number of host cells transformed with a nucleic acid of the invention  
will depend, at least in part, upon the type of recombinant expression vector used and the  
type of transfection technique used. Nucleic acid can be introduced into a host cell  
30 transiently, or for long term. In long-term systems, the nucleic acid is stably integrated into  
the genome of the host cell or remains as a stable episome in the host cell.

Certain vectors integrated into host cells at only a low frequency. In order to  
identify these integrants, a gene that contains a selectable marker (*e.g.*, drug resistance) is  
generally introduced into the host cells along with the nucleic acid of interest. Preferred  
35 selectable markers include those which confer resistance to certain drugs, such as G418 and  
hygromycin. Selectable markers can be introduced on a separate plasmid from the nucleic

5 acid of interest or, are introduced on the same plasmid. Host cells transfected with a nucleic acid of the invention (e.g., a recombinant expression vector) and a gene for a selectable marker can be identified by selecting for cells using the selectable marker. For example, if the selectable marker encodes a gene conferring neomycin resistance, host cells which have taken up nucleic acid can be selected with G418. Cells that have incorporated the selectable  
10 marker gene will survive, while the other cells die.

Nucleic acid encoding the polypeptides of the invention can be introduced into cells growing in culture *in vitro* by conventional transfection techniques (e.g., calcium phosphate precipitation, DEAE-dextran transfection, electroporation, biolistics, etc.). Nucleic acid can also be transferred into cells *in vivo*, for example by application of a  
15 delivery mechanism suitable for introduction of nucleic acid into cells *in vivo*, such as retroviral vectors (see e.g., Ferry *et al.* (1991) *Proc. Natl. Acad. Sci., USA*, 88: 8377-8381; and Kay *et al.* (1992) *Human Gene Therapy* 3: 641-647), adenoviral vectors (see, e.g., Rosenfeld (1992) *Cell* 68: 143-155; and Herz and Gerard (1993) *Proc. Natl. Acad. Sci., USA*, 90:2812-2816), receptor-mediated DNA uptake (see e.g., Wu, and Wu (1988) *J. Biol. Chem.* 263: 14621; Wilson *et al.* (1992) *J. Biol. Chem.* 267: 963-967; and U.S. Pat. No. 5,166,320), direct injection of DNA (see, e.g., Acsadi *et al.* (1991) *Nature* 332: 815-818; and Wolff *et al.* (1990) *Science* 247:1465-1468) or particle bombardment (biolistics) (see e.g., Cheng *et al.* (1993) *Proc. Natl. Acad. Sci., USA*, 90:4455-4459; and Zelenin *et al.* (1993) *FEBS Letts.* 315: 29-32).

25 **5) Recovery of expressed polypeptide or holophytochrome.**

In some instances, it is desired to recover the expressed polypeptide (e.g. HY2 family reductase) and/or the assembled holophytochrome or the holophytochrome labeled polypeptide. Once expressed, the desired proteins and/or holophytochromes can be purified according to standard procedures of the art, including ammonium sulfate  
30 precipitation, affinity columns, column chromatography, gel electrophoresis and the like (see, generally, R. Scopes, (1982) *Protein Purification*, Springer-Verlag, N.Y.; Deutscher (1990) *Methods in Enzymology Vol. 182: Guide to Protein Purification.*, Academic Press, Inc. N.Y.). In certain embodiments, substantially pure compositions of at least about 90 to 95% homogeneity are preferred, and 98 to 99% or more homogeneity are most preferred.  
35 The cloning and expression of a HY2 family members is illustrated in Examples 1 and 2, and the expression of a holophytochrome is illustrated in Example 3.

5 One of skill would recognize that modifications can be made to the apophytochrome and/or the HY2 reductases (or other components of the phytochrome synthetic pathway) without diminishing their biological activity. Some modifications may be made to facilitate the cloning, and expression of the subject molecule(s). Such modifications are well known to those of skill in the art and include, for example, a methionine added at the amino terminus to provide an initiation site, or additional amino acids (*e.g.*, poly His) placed on either terminus to create conveniently located restriction sites or termination codons or purification sequences.

#### **IV. Assembly of phytochromes and phytofluors.**

15 In certain preferred embodiments, this invention provides for the assembly of holophytochromes. It was a surprising discovery of this invention that a cell transfected with nucleic acids encoding the components of a bilin synthetic pathway (*e.g.*, HO1, PcyA, and/or HY2) and a nucleic acid encoding an apophytochrome (*e.g.* Cph1)) will express a phytochromobilin that assembles with the apophytochrome to produce a photoactive holophytochrome.

20 It has been demonstrated that recombinant apophytochromes produced in microorganisms can self assemble with the bilins, phycocyanobilin, phytochromobilin and phycoerythrobilin, to produce photoreversible holophytochromes and intensely fluorescent phytofluors *in vitro* (Wahleithner *et al.* (1991) *Proc. Natl. Acad. Sci. USA*, 88: 10387-10391; Li and Lagarias (1992) *J. Biol. Chem.*, 267: 19204-19210; Murphy and Lagarias (1997) *Curr. Biol.* 7: 870-876; US Patent 6,046,041).

This invention additionally provides the genes encoding ferredoxin-dependent bilin reductases that convert biliverdin to phytochromobilin, phycocyanobilin or phycoerythrobilin (see Figure 1). In one aspect, this invention describes an *in vivo* expression system for holophytochrome.

30 One preferred approach involved the design of a synthetic operon comprising HO1 and PcyA coding regions from *Synechocystis* sp. PCC6803 (Yanofsky *et al.* (1981) *Nucl. Acids Res.*, 9: 6647-6667; Baneyx (1999) *Curr. Opin. Biotechnology*, 10: 411-421; Yeh, *et al.* (1997) *Science*, 277: 1505-1508). Cloning of *ho1* and *pcyA* genes from *Synechocystis* sp. PCC6803 in the plasmid pPROLarA122 (Clontech Laboratories) places these genes under the control of dual *Ara/Lac* promotor. Upon introduction of this plasmid

35

5 into *E. coli* cells harboring the Cph1-expression plasmid, pBAD/Cph1(514), photoactive holophytochrome is expressed/assembled *in vivo*.

Another particularly preferred approach, illustrated in Example 3, involved the production of a synthetic operon comprised of *HO1* from *Synechocystis* sp. PCC6803 and the mature *HY2* coding region (mHY2) from *Arabidopsis thaliana* that lacks the plastid targeting sequence. The cloning of *HO1* and mHY2 open reading frames into the plasmid pPROLarA122 (Clontech Laboratories) placed this operon under regulatory control of a dual *Ara/Lac* promoter. Upon introduction of this plasmid into *E. coli* cells harboring the Cph1-expression plasmid, pBAD/Cph1(514), in which Cph1(N514) is under regulatory control of a *Ara* promoter, the production of photoactive holophytochrome *in vivo* was  
15 observed.

These approaches are illustrative and not meant to be limiting. Using the teaching provided herein, numerous other approaches will be available to one of skill in the art. For example, cell lines naturally harboring HO1 can be used thereby eliminating the need to provide this enzyme from a heterologous nucleic acid. Alternatively, the cell can be  
20 provided exogenous biliverdin using a variety of transfection reagents (*e.g.* (*e.g.* cationic lipids, lipofectamine<sup>TM</sup>, Chariot<sup>TM</sup>, etc.).

Other bilin reductases can be used, numerous apoproteins or minimal domains thereof sufficient to form holophytochromes and/or phytofluors can be used, and other components of the bilin biosynthetic pathway can be provided by heterologous nucleic acids. For example, it is cells expressing an apophytochrome, HO1, *pebA* and *pebB* will produce phytofluors *in vivo*. Similarly, co-expression of the structural gene for a phycobiliprotein, a phycobiliprotein bilin lyase and the genes necessary for a phytyobilin biosynthetic pathway from heme will lead to the production of fluorescent phycobiliproteins in living cells.

## 30 V. Kits.

This invention also provides kits for the practice of the methods of this invention. In one embodiment the kits include a container containing one or more bilin reductases of this invention (*e.g.* HY2 family members) and/or nucleic acids encoding one or more bilin reductases of this invention. In certain embodiments, the kits comprise a  
35 container containing nucleic acids sufficient express and assemble a holophytochrome (*e.g.* a bilin chromophore or a phytofluor) in a host cell. Such kits, optionally include a vector

5 encoding an apoprotein and, optionally, a restriction site to insert a nucleic acid into the vector so the heterologous nucleic acid expresses a fusion protein with the apoprotein.

The kits may optionally include devices and reagents to facilitate performing the methods of this invention. Such devices and reagents include, but are not limited to microtiter plates (*e.g.* for high-throughput applications), culture plates, culture media, cell  
10 lines, buffers, labels, and the like.

In addition, the kits optionally include labeling and/or instructional materials providing directions (*i.e.*, protocols) for the practice of the methods of this invention. Preferred instructional materials describe the expression of a bilin reductase and/or the *in vivo* expression/assembly of a holophytochrome and/or a phytofluor and/or the expression  
15 of a polypeptide labeled (as a fusion protein) with a holophytochrome or a phytofluor.

While the instructional materials typically comprise written or printed materials they are not limited to such. Any medium capable of storing such instructions and communicating them to an end user is contemplated by this invention. Such media include, but are not limited to electronic storage media (*e.g.*, magnetic discs, tapes, cartridges,  
20 chips), optical media (*e.g.*, CD ROM), and the like. Such media may include addresses to internet sites that provide such instructional materials.

### EXAMPLES

The following examples are offered to illustrate, but not to limit the claimed invention.

25

#### Example 1

#### The Arabidopsis HY2 Gene Encodes Phytochromobilin Synthase, a Ferredoxin-Dependent Biliverdin Reductase

Light perception by the plant photoreceptor phytochrome requires the tetrapyrrole chromophore phytochromobilin (PΦB), which is covalently attached to a large  
30 apoprotein. Arabidopsis mutants *hy1* and *hy2*, which are defective in PΦB biosynthesis, display altered responses to light due to a deficiency in photoactive phytochrome. In this example, we describe the isolation of the *HY2* gene by map-based cloning. *hy2* mutant alleles possess alterations within this locus, some of which affect the expression of the *HY2* transcript. *HY2* encodes a soluble protein precursor of 38 kD with a putative N-terminal  
35 plastid transit peptide. The *HY2* transit peptide is sufficient to localize the reporter green

5 fluorescent protein to plastids. Purified mature recombinant HY2 protein exhibits PΦB synthase activity (*i.e.*, ferredoxin-dependent reduction of biliverdin IX $\alpha$  to PΦB), as confirmed by HPLC and by the ability of the bilin reaction products to combine with apophytochrome to yield photoactive holophytochrome. Database searches and hybridization studies suggest that *HY2* is a unique gene in the Arabidopsis genome that is  
10 related to a family of proteins found in oxygenic photosynthetic bacteria.

### **Introduction.**

Plants are exquisitely sensitive to their environment. Because they are sessile and use light as the energy source for photosynthesis, plants have developed well-refined photoreception and signaling systems to modulate their growth and development.  
15 The family of phytochromes, which are sensory photoreceptors for red and far red light, play a key role in mediating responses to light quality, quantity, direction, and duration throughout plant development (Kendrick and Kronenberg (1994) *Photomorphogenesis in Plants*. (Dordrecht, The Netherlands: Martinus Nijhoff Publishers); Quail *et al.* (1995) *Science* 268: 675-680; Furuya, and Schäfer (1996) *Trends Plant Sci.* 1: 301-307; Neff *et al.* (2000) *Genes Dev.* 14: 257-271). Plant phytochromes are homodimers composed of ~125-  
20 kD subunits each with a thioether-linked phytochromobilin (PΦB) prosthetic group (Lagarias and Rapoport (1980) *J. Am. Chem. Soc.* 102: 4821-4828). Phytochrome action depends on its ability to photointerconvert between the red light-absorbing form and the far-red-light absorbing form, a property conferred by covalently bound PΦB in  
25 holophytochrome.

Two pathways are involved in the biosynthesis of holophytochrome, one for the apoprotein, which is encoded by a small multigene family (*e.g.*, *PHYA-E* in Arabidopsis) (Sharrock and Quail (1989) *Genes Dev.* 3: 1745-1757; Clack *et al.* (1994) *Plant Mol. Biol.* 25: 413-427), and another for the synthesis of the PΦB (Terry *et al.* (1993) *Arch. Biochem. Biophys.* 306: 1-15). Apophytochrome is synthesized in the cytosol,  
30 whereas PΦB is synthesized entirely within the plastid compartment, followed by its release to the cytosol, where holophytochrome assembly occurs (Figure 2). Based on spectroscopic studies of purified phytochromes, in vitro bilin assembly studies with recombinant apophytochromes, and physiological analyses of chromophore-deficient mutants, PΦB  
35 appears to be the immediate chromophore precursor of all higher plant and cryptophyte

5 phytochromes (Terry *et al.* (1993) *Arch. Biochem. Biophys.* 306: 1-15; Terry (1997) *Plant Cell Environ.* 20: 740-745).

PΦB is synthesized from 5-aminolevulinic acid and shares many intermediates with the pathways of chlorophyll and heme biosynthesis (Elich and Lagarias (1987) *Plant Physiol.* 84: 304-310; Elich *et al.* (1989) *J. Biol. Chem.* 264: 183-189). These analyses established that biliverdin IXα (BV) is a PΦB precursor, suggesting the intermediacy of heme in the phytochrome chromophore biosynthetic pathway. Indeed, the first committed step of PΦB biosynthesis is catalyzed by a ferredoxin-dependent heme oxygenase, which is encoded by the *HY1* gene in Arabidopsis and by its ortholog in rice (Davis *et al.* (1999) *Proc. Natl. Acad. Sci., USA*, 96: 6541-6546; Muramoto *et al.* (1999) *Plant Cell* 11: 335-347; Izawa *et al.* (2000) *Plant J.* 22: 391-399). Ferredoxin-dependent heme oxygenases were first identified in red algae and cyanobacteria, in which they catalyze the oxygen-dependent conversion of heme to BV (Beale and Cornejo (1984) *Arch. Biochem. Biophys.* 235, 371-384; Cornejo and Beale (1988) *J. Biol. Chem.* 263: 11915-11921; Cornejo and Beale (1997) *Photosynth. Res.* 51: 223-230; Cornejo *et al.* (1998) *Plant J.* 15: 99-107). BV, therefore, is the first committed intermediate in the biosynthetic pathways of PΦB as well as those of the phycobilins phycocyanobilin and phycoerythrobilin, which are precursors of the light-harvesting prosthetic groups of the phycobiliproteins in cyanobacteria, red algae, and cryptomonads (Beale (1993) *Chem. Rev.* 93: 785-802).

25 In plants, BV is subsequently reduced to 3Z-PΦB by the ferredoxin-dependent bilin reductase PΦB synthase, which has not yet been cloned (Terry and Lagarias (1991) *J. Biol. Chem.* 266: 22215-22221). Although 3Z-PΦB can serve as a functional precursor of the phytochrome chromophore, its facile isomerization to 3E-PΦB, which is also a precursor of the phytochrome chromophore, likely occurs in plants (Terry *et al.* (1995) *J. Biol. Chem.* 270: 11111-11118). Ferredoxin-dependent bilin reductases are also present in cyanobacteria and red algae, where they catalyze the conversion of BV to the phycobilins (reviewed by Beale (1993) *Chem. Rev.* 93: 785-802). None of these bilin reductases has previously been cloned.

35 Our understanding of photomorphogenesis in plants has been aided greatly by the isolation of five classic photomorphogenic Arabidopsis mutants (*hyl* to *hy5*) that are impaired in response to light (Koornneef *et al.* (1980) *Z. Pflanzenphysiol.* 100: 147-160).

- 5 Photoreceptor-deficient mutants have proven to be powerful tools to analyze which photoreceptors mediate specific photomorphogenetic responses (Koornneef and Kendrick (1994) *Photomorphogenic mutants of higher plants*. Pp. 601-628 In *Photomorphogenesis in Plants*, R.E. Kendrick and G.H.M. Kronenberg, eds Dordrecht, The Netherlands: Kluwer Academic Pub.; Whitelam and Devlin (1997) *Plant Cell Environ.* 20: 752-758).
- 10 Phytochrome chromophore-deficient mutants, including *hyl* and *hy2* in *Arabidopsis*, *yg-2* and *aurea* in tomato, *pcd1* and *pcd2* in pea, and *pew1* and *pew2* in *Nicotiana glumbaginifolia*, have often been used as phytochrome-deficient mutants (reviewed by Terry (1997) *Plant Cell Environ.* 20: 740-745). The *aurea* mutant of tomato has been used widely for physiological studies of phytochrome, for the study of other photoreceptors, and
- 15 to study phytochrome signaling (Becker *et al.* (1992) *Planta* 188: 39-47; Bowler and Chua (1994) *Plant Cell* 6: 1529-1541). Knowledge of the molecular basis of these mutations will help in the interpretation of physiological experiments with these mutants. Biochemical analyses have established that the *hyl*, *pcd1*, and *yg-2* mutants are deficient at the step at which BV is synthesized from heme, whereas *pcd2* and *aurea* mutants are unable to
- 20 synthesize PΦB from BV (Terry and Kendrick (1996) *J. Biol. Chem.* 271: 21681-21686; van Tuinen *et al.* (1996) *Plant J.* 9: 173-182; Weller *et al.* (1996) *Plant Cell* 8: 55-67; Weller *et al.* (1997) *Plant J.* 11: 1177-1186). The cloning of *HY1* has provided valuable insight into the first committed enzyme of phytochrome chromophore biosynthesis, heme oxygenase (Davis *et al.* (1999) *Proc. Natl. Acad. Sci., USA*, 96: 6541-6546; Muramoto *et al.* (1999) *Plant Cell* 11: 335-347).
- 25

Of the five classic photomorphogenetic mutants, only *hy2* remains to be cloned. It is widely believed that *HY2* encodes PΦB synthase. However, the observation that a *hy2* mutant is partially "rescued" by BV treatment suggests other possibilities (Parks and Quail (1991) *Plant Cell* 3: 1177-1186). Although it is similar to *hyl* mutants, the

30 chlorophyll-deficient phenotype of *hy2* mutants is typically less severe (Koornneef *et al.* (1980) *Z. Pflanzenphysiol.* 100: 147-160; Chory *et al.* (1989) *Plant Cell* 1: 867-880). The gene identification of *HY2* in *Arabidopsis* should help to resolve these paradoxes. In this study, we describe the molecular basis for the phytochrome-deficient phenotype in the *hy2* mutant of *Arabidopsis*. We show that the *HY2* gene encodes PΦB synthase, a ferredoxin-dependent BV reductase that is responsible for the final step in phytochrome chromophore

35 biosynthesis in plastids. This work has enabled us to identify other members of the *HY2*-



- 5 related, ferredoxin-dependent bilin reductase family in phycobiliprotein-producing photosynthetic organisms (see Example 3 herein).

### Results.

#### Fine Mapping Localizes the *HY2* Gene to Two Overlapping Bacterial Artificial Chromosome Clones

- 10 We used a positional cloning strategy to isolate the *HY2* gene, which previously had been mapped to chromosome 3. Because the *hy2* long hypocotyl phenotype is easy to score in seedlings, the *HY2* locus has served as a useful landmark for classic mapping. For fine mapping, we crossed the *hy2-1* mutant of Landsberg *erecta* (*Ler*) ecotype to the wild-type Columbia ecotype, and segregating F2 populations with the *hy2* phenotype were used for DNA preparation. First, we prepared DNA from ~400 plants to perform genetic mapping of *hy2* using cleaved amplified polymorphic sequence (CAPS) markers (Konieczny and Ausubel (1993) *Plant J.* 4: 403-410) that we developed and that are available in the database at the Arabidopsis Information Resource (TAIR; [http://www.arabidopsis.org/maps/CAPS\\_Ch3.html](http://www.arabidopsis.org/maps/CAPS_Ch3.html)). With ~400 plants, *HY2* was mapped to an interval of ~360 kb between positional markers cMLP3E-1 and cF3L24 (Figure 3A), indicating that recombination frequency in this region was much lower than expected. Therefore, we increased the size of the mapping population to ~2000 plants. This approach enabled us to map the *HY2* locus to an interval of ~66 kb between the markers cMZB10 and cF3L24 (Figure 3A).

- 25 During these mapping studies, the sequences of two bacterial artificial chromosome clones, MZB10 and F3L24, spanning the *HY2* locus genetically defined above, were deposited in the GenBank database (accession numbers AC009326 and AC011436). There are at least 21 putative genes in the region between the closest recombinations. We screened *HY2* candidate genes based on the following expectations. First, *HY2* should be categorized as an unknown or putative gene, because neither gene nor protein sequences of any ferredoxin-dependent bilin reductase were known. Second, *HY2* should possess a plastid transit peptide, because enzymatic activity for PΦB synthase was detected in plastids (Terry and Lagarias (1991) *J. Biol. Chem.* 266: 22215-22221). Third, weak sequence similarity between *HY2* and an unidentified open reading frame(s) (ORFs) in fully sequenced cyanobacterial bacterial genomes might be detectable, because *HY2*-related bilin
- 30
- 35

5 reductase activities have been reported in cyanobacteria (Cornejo and Beale (1997) *Photosynth. Res.* 51: 223-230). The predicted amino acid sequences for all 21 genes in the *HY2* region were used for TBLASTN (Altschul *et al.* (1990) *J. Mol. Biol.* 215: 403-410) and CHLOROP (Emanuelsson *et al.* (1999) *Protein Sci.* 8: 978-984; <http://www.cbs.dtu.dk/services/ChloroP/>) analyses. By these criteria, one of these genes  
10 with two distinct annotations, MZB10.18 (GenBank accession number AC009326-18) or F3L24.1 (GenBank accession number AC0011436-1), appeared to be a strong candidate for *HY2*.

**The *HY2* Gene Is Identified by DNA Sequences of Wild-Type and Mutant Alleles**

15 To help identify the *HY2* gene, RNA gel blot analysis of wild-type and *hy2* mutant seedlings was performed using the cDNA for MZB10.18/F3L24.1 as a probe. Because the *hy2* phenotype is readily observed in seedlings, we analyzed the accumulation of transcripts in Arabidopsis seedlings (Figure 4). Transcripts were detected in wild-type of three ecotypes tested. The slow migration of mRNA of Col was verified as a gel artifact  
20 (data not shown). RNA gel blotting showed that the transcript levels were decreased severely in the *hy2-1*, *hy2-106*, and *hy2-107* mutants and were decreased slightly in other mutant lines. Consequently, we focused our attention on the MZB10.18/F3L24.1 gene. To determine if mutations were present in the MZB10.18/F3L24.1 gene in *hy2* mutants, DNA fragments corresponding to the region from the end of the upstream gene to the beginning  
25 of the downstream gene from various *hy2* alleles were amplified by polymerase chain reaction (PCR). The nucleotide sequences were determined directly from the PCR products. In all *hy2* alleles tested, nucleotide substitutions or deletions were detected (Figure 3A and 3B). Based on these data and biochemical data presented below, we conclude that locus MZB10.18/F3L24.1 corresponds to the *HY2* gene.

30 As a result of the conflict in annotation of the *HY2* gene in MZB10.18 and F3L24.1 (*i.e.*, the former encodes a protein of 273 amino acids, and the latter encodes a protein of 329 amino acids), we sought to verify experimentally the structure of the *HY2* gene. To do so, seven cDNA clones prepared from Columbia seedling mRNA were isolated from ~300,000 clones examined. The nucleotide sequences of independent cDNA clones  
35 were determined, and they revealed a single reading frame that matched that of the

5 annotation for F3L24.1. The *HY2* gene contains eight small exons ranging from 51 to 222 nucleotides separated by seven introns ranging from 74 to 183 nucleotides. The longest cDNA insert contained a full length 990-bp ORF, a 95-bp 5'-untranslated region, a 231-bp 3'-untranslated region, and a poly(A)<sup>+</sup> stretch, as shown in Figure 5 (DNA Data Bank of Japan [DDBJ] accession number AB045112).

10 Figure 3A shows the genomic structure of the *HY2* gene with positions of the mutations in *hy2* alleles. Two *hy2* alleles, *hy2-102* and *hy2-107*, were found to have point mutations at 3' splice sites in the seventh and fifth introns, respectively. Such mutations in the G of the essential AG dinucleotide at the 3' splice site have been reported to lead to missplicing with a downstream AG, resulting in a frameshift in the protein (Brown (1996) *Plant J.* 10: 771-780). *hy2-105* was another possible splicing mutant, with a 25-bp deletion in the second intron. This mutation truncates the second intron to 57 nucleotides, much smaller than the average size of Arabidopsis introns (240 nucleotides). The efficiency of intron splicing might be reduced because of a minimum intron size requirement (Deutsch and Long (1999) *Nucleic Acids Res.* 27: 3219-3228), although we have not checked the significance of defects in pre-mRNA splicing experimentally. A fast neutron-generated allele, *hy2-106*, carries a 5-bp deletion in the first exon, making an immediate stop codon. Four ethyl methanesulfonate-generated alleles, *hy2-1*, *hy2-101*, *hy2-103*, and *hy2-104*, have single nucleotide changes to produce amino acid substitutions compared with the corresponding wild-type allele. Two of these alleles, *hy2-1* and *hy2-104*, have the same mutation (P128L), whereas *hy2-101* and *hy2-103* possess G181R and R252Q substitutions, respectively.

### **The HY2 Protein Is Related to a Family of Cyanobacterial Proteins.**

Based on cDNA sequence analysis, the *HY2* protein contains 329 residues with a calculated molecular mass of 38.1 kD. At its N terminus, the *HY2* protein sequence is rich in serine, with few acidic residues (six serine and one aspartic acid among 45 residues), which suggests a possible transit peptide for localization to plastids (Gravel and von Heijne (1990) *FEBS Lett.* 261: 455-458). The second amino acid after the initiation methionine is alanine, which is often observed in plastid transit peptides. The program CHLOROP was also used to predict the transit peptide of *HY2*, and it indicated that the first 45 amino acid residues of the *HY2* protein form a chloroplast transit peptide (Emanuelsson

5 *et al.* (1999) *Protein Sci.* 8: 978-984; <http://www.cbs.dtu.dk/services/ChloroP/>). The calculated molecular mass of the mature HY2 protein is 33.0 kD and its predicted pI is 5.66, which are similar to those of PΦB synthase purified from oat seedlings. The HY2 protein has no predicted transmembrane helices, which is also consistent with the observation that oat PΦB synthase is a soluble protein.

10 Using the HY2 protein sequence as a query sequence, we performed an iterative PSI-BLAST search of the nonredundant GenBank/EMBL database (<http://www.ncbi.nlm.nih.gov/blast/psiblast.cgi>) using default search parameters (Altschul *et al.* (1997) *Nucleic Acids Res.* 25, 3389-3402). Surprisingly, no HY2-related gene was identified by this search in the nearly complete Arabidopsis genome. In contrast, this search  
15 identified HY2-related sequences from two marine cyanobacteria, *Prochlorococcus marinus* sp. SS120 (EMBL accession numbers CAB95700.1 and CAB95701.1) and *Synechococcus* sp. WH8020 (SWISS-PROT accession numbers Q02189 and Q02190), and a related protein sequence from the cyanobacterium *Synechocystis* sp. PCC 6803 (cyanobase locus *slr0116*; Protein Information Resource accession number S76709). Both marine cyanobacteria  
20 possess two HY2-related ORFs that appear to be part of multigene operons. Interestingly, the *Synechococcus* ORFs, *ycp2\_synpy* and *ycp3\_synpy*, are located within a cluster of genes involved in phycobiliprotein biosynthesis (Wilbanks and Glazer (1993) *J. Biol. Chem.* 268: 1226-1235), whereas the *Prochlorococcus* ORFs, which we term *yhp2\_proma* and *yhp3\_proma*, are located immediately downstream of a gene related to heme oxygenase  
25 (GB:AJ278499.1). These observations strongly support the hypothesis that these genes are involved in phycobilin biosynthesis.

Figure 5 shows an optimized multiple sequence alignment of HY2 and HY2-related cyanobacterial sequences using the programs CLUSTALW (Higgins *et al.* (1996) *Meth. Enzymol.* 266: 383-402), MEME to guide hand alignments  
30 (<http://meme.sdsc.edu/meme/website/>), and GENEDOC for highlighting (<http://www.psc.edu/biomed/genedoc>). As expected, the HY2-related cyanobacterial proteins lack the putative plastid transit peptide sequence found at the N terminus of HY2. Pairwise sequence identities between HY2 and the cyanobacterial ORFs are quite low (<20%), although the similarities between YCP2\_SYNPY and YHP2\_PROMA and  
35 between YCP3\_SYNPY and YHP3\_PROMA suggest that these pairs of proteins have similar functions. That the mutation in the *hy2-1* and *hy2-104* alleles (P128L) lies in a

5 conserved proline residue is consistent with a critical role of this residue in the enzyme's structure. Proline residues are typically involved in *cis*-peptide bonds, which occur at  $\beta$ -turns in proteins. Examination of the amino acid alterations in the two other missense alleles, G181R in *hy2-101* and R252Q in *hy2-103*, reveals that neither mutation corresponds to a strongly conserved residue in this protein family.

#### 10 **The HY2 Protein Is Localized to the Plastid.**

The N terminus of HY2 has a stretch of 45 amino acids with features of a chloroplast transit peptide. To determine whether this peptide is a functional plastid-targeting sequence, we fused the transit peptide coding region of *HY2* to a modified gene of green fluorescent protein (GFP) from jellyfish under the control of modified cauliflower  
15 mosaic virus 35S promoter (Chiu *et al.* (1996) *Curr. Biol.* 6: 325-330). The construct was introduced into onion skin cells and tobacco leaves by bombardment with DNA-coated particles, and transient expression was analyzed using confocal laser scanning microscopy. Although a control construct without the putative transit peptide showed GFP fluorescence throughout the cytoplasm and the nucleus of onion cells (Figure 6A), clear localization of  
20 GFP fluorescence to small dots, most likely plastids, was observed when the putative transit peptide was fused to GFP (Figure 6B). For better visualization, we also introduced the construct into tobacco leaves, where the chloroplasts are well developed in guard cells. GFP fluorescence was localized exclusively in oval structures (Figure 6C) that match the red autofluorescence from the chlorophyll of the chloroplasts (Figure 6D), demonstrating that  
25 the fusion protein is efficiently targeted to chloroplasts. This finding confirms the presence of a functional transit peptide and implies that the *HY2* gene product is localized in the chloroplast.

#### **Recombinant HY2 Exhibits P $\Phi$ B Synthase Activity**

The HY2 protein lacking the transit peptide, mHY2, was synthesized in  
30 *Escherichia* as a fusion protein with glutathione-S-transferase (GST) and purified by affinity chromatography, as described in Methods. The GST tag was removed by site-specific protease digestion. A second round of affinity chromatography yielded protein at >90% homogeneity. Figure 7 shows SDS-PAGE results of the purification and processing of the protein. One liter of bacterial culture yielded approximately 1 mg of recombinant protein.  
35 The molecular mass of the Arabidopsis mHY2 deduced from the cDNA is 33 kD.

- 5 However, the cloning and expression strategy for the mHY2 cDNA using pGEX-6P-1 was responsible for an additional five N-terminal amino acids (GPLGS) after protease treatment.

To determine whether mHY2 has P $\Phi$ B synthase activity, its ability to reduce BV to P $\Phi$ B was first assessed with a "coupled" holophytochrome assembly assay in which the reaction products were incubated with recombinant cyanobacterial phytochrome 1  
10 (Cph1) apoprotein (Yeh *et al.* (1997) *Science* 277: 1505-1508).

Figure 8 shows a phytochrome difference spectrum obtained after incubation of apoCph1 with the bilin products from a P $\Phi$ B synthase assay of a crude cell-free bacterial extract expressing GST-mHY2. The difference spectrum has a peak at 676 nm and a valley at 724 nm, which is consistent with a P $\Phi$ B-Cph1 adduct (Yeh *et al.* (1997) *Science* 277:  
15 1505-1508). To ensure that this activity was not due to a component of the crude *Escherichia* lysate, the ability of purified mHY2 to reduce BV to P $\Phi$ B was analyzed using the coupled assembly assay as well as an HPLC assay. A phytochrome difference spectrum identical to that shown in Figure 8 was obtained (data not shown). The HPLC results of the P $\Phi$ B synthase assay mixture are shown in Figure 9. After incubation of HY2 for 30 min  
20 under standard P $\Phi$ B synthase assay conditions, all of the BV was converted to P $\Phi$ B. Interestingly, both 3Z- and 3E-P $\Phi$ B isomers were produced, although the relative amount of the 3E-P $\Phi$ B isomer varied for different HY2 samples and may be an artifact of the presence of residual glutathione.

### Discussion.

- 25 The *hy2* mutant of *Arabidopsis* is one of five classic long hypocotyl mutants first identified by Koornneef *et al.* (1980) *Z. Pflanzenphysiol.* 100: 147-160. That the *hy2* mutant is photomorphogenetically impaired due to a phytochrome deficiency has been well documented by physiological and photobiological analyses (Koornneef *et al.* (1980) *Z. Pflanzenphysiol.* 100: 147-160; Chory *et al.* (1989) *Plant Cell* 1: 867-880; Parks and Quail  
30 (1991) *Plant Cell* 3: 1177-1186; Goto *et al.* (1993) *Photochem. Photobiol.* 57: 867-871). Parks and Quail (1991) *Plant Cell* 3: 1177-1186, showed that the long hypocotyl phenotype of the *hy1* and *hy2* mutants was in part "rescued" by BV feeding and suggested that these mutants have lesions in the phytochrome chromophore biosynthetic pathway. Indeed, *HY1* encodes a plastid-localized heme oxygenase that catalyzes the cleavage of heme to form BV  
35 (Davis *et al.* (1999) *Proc. Natl. Acad. Sci., USA*, 96: 6541-6546; Muramoto *et al.* (1999)

5 *Plant Cell* 11: 335-347). This example establishes that *HY2* encodes PΦB synthase, a plastid-localized enzyme responsible for the ferredoxin-dependent conversion of BV to PΦB, the immediate precursor of the phytochrome chromophore. Although complementation experiments are in progress, sequence analysis of eight mutant alleles has revealed molecular lesions within the *HY2* gene. Many of the *hy2* alleles also display  
10 altered expression of the *HY2* transcript, providing compelling evidence that the reduced expression of this gene is responsible for the long hypocotyl phenotype.

Based on the presence of a functional plastid-targeting sequence in the *HY2* protein, we can confidently conclude that the entire pathway of PΦB biosynthesis occurs within plastids. Nevertheless, the possibility of an alternative pathway in other subcellular  
15 compartments cannot be dismissed entirely. In this regard, there are three other heme oxygenase genes besides *HY1* in the Arabidopsis genome whose products may play a role in an alternative pathway (M. Masuda, T. Muramoto, and T. Kohchi, unpublished data). However, our database searches revealed no other gene in the Arabidopsis genome that shows statistically significant similarity to *HY2*. Although a weak similarity between *HY2*  
20 and a ferredoxin-dependent bilin reductase involved in chlorophyll catabolism, red chlorophyll catabolite reductase, was revealed by profile analysis, red chlorophyll catabolite reductase does not catalyze the reduction of BV to PΦB (Wüthrich *et al.* (2000) *Plant J.* 21: 189-198). Therefore, it appears that *HY2* is the only PΦB synthase gene in Arabidopsis.

Physiological comparisons of the *hy1* and *hy2* mutants indicate that *hy1*  
25 plants display more severe phytochrome-deficient phenotypes (Koornneef *et al.* (1980) *L. Heynh. Z. Pflanzenphysiol.* 100: 147-160; Chory (1989) *Plant Cell* 1: 867-880). These observations are somewhat surprising in view of the apparent uniqueness of the *HY2* gene and the existence of multiple *HY1*-related proteins in the Arabidopsis genome. However, this may reflect the strength of the *hy1* and *hy2* alleles examined. In this regard, the partial  
30 rescue of the *hy2-1* mutant treated with BV (Parks and Quail (1991) *Plant Cell* 3: 1177-1186) can be explained by the hypothesis that the P131L missense mutation affords a partially active enzyme with a lower affinity for BV. Alternatively, it is possible that BV might be converted to PΦB by an enzyme unrelated to *HY2* in Arabidopsis. Phytochrome chromophore biosynthetic mutants have been identified in other plant species (Terry (1997)  
35 *Plant Cell Environ.* 20: 740-745). In all cases, two classes of mutants have been identified: those that are deficient in heme oxygenase and those that are deficient in PΦB synthase.

5 Based on biochemical analyses, the *aurea* mutant of tomato and the *pcd2* mutant of pea are deficient in PΦB synthase activity (van Tuinen *et al.* (1996) *Plant J.* 9: 173-182; Weller *et al.* (1997) *Plant J.* 11: 1177-1186). The observations that the corresponding heme oxygenase mutants in these plant species (*i.e.*, *yg-2* and *pcd1*, respectively) exhibit less severe phenotypes further support the hypothesis that the relative allele strength of the two  
10 loci determines the phenotype. A phenotypic comparison of null alleles of *hyl* and *hy2* (*e.g.*, *hy2-106* and *hy2-107*) should help resolve this question.

The cloning of the Arabidopsis *HY2* gene will help to identify PΦB synthase genes from other plant species and to confirm that the mutations in *aurea* and *pcd2* occur in homologous genes. The *aurea* mutant of tomato has been used extensively to analyze  
15 phytochrome signal transduction (Bowler *et al.* (1994) *Cell* 77: 73-81), and knowledge of the molecular basis of this mutation is of considerable interest. The molecular basis of such mutations should provide insight into residues critical for substrate and/or potential cofactor (*i.e.*, metal ions or organic single electron carriers) interactions as well as those necessary for protein-protein interactions (*i.e.*, between *HY2* and ferredoxin or between *HY1* and  
20 *HY2*). The availability of *HY1*- and *HY2*-specific cDNA probes and specific antibodies to both enzymes will facilitate experiments to study the regulation of phytochrome chromophore biosynthesis. With such probes, several key questions can be addressed. Are the two enzymes expressed coordinately in all tissues? Is their expression spatially and temporally regulated? Do *HY1* and *HY2* proteins form a dual enzyme complex in the  
25 plastid that channels the conversion of heme to PΦB? Does the expression of *HY1* affect *HY2* expression and vice versa?

The molecular cloning of *HY2* has provided a breakthrough in our knowledge of bilin biosynthesis in general. Our bioinformatic analyses reveal that *HY2* is related to a number of cyanobacterial genes of unknown function (Figure 5). Indeed, we  
30 believe these *HY2*-related proteins are enzymes involved in the biosynthesis of the chromophore precursors of the light-harvesting phycobiliproteins phycocyanobilin and phycoerythrobilin. As might be expected for enzymes with different substrate/product specificities, these proteins are highly diverged from *HY2* (<20% sequence identity). The levels of identity between these proteins and *HY2*, which are highlighted in Figure 5, likely  
35 reflect residues involved in overall protein folding and/or ferredoxin interaction that are common to the entire family of enzymes. In Example 2, we demonstrate that these *HY2*-



5 related proteins are members of a growing family of ferredoxin-dependent bilin reductases with different double bond specificities.

The pathway for phytochrome chromophore biosynthesis shown in Figure 1 has been clearly documented. Now that the two key genes of the phytochrome chromophore biosynthetic pathway have been cloned, we can elucidate how bilin  
10 biosynthesis is regulated throughout the plant, a process that is critical to the plant's ability to respond to light. The possible role of bilins as second messengers, which was raised by recent studies of transgenic plants expressing mammalian biliverdin reductase (Montgomery *et al.* (1999) *Plant Physiol.* 121: 629-639), can be addressed by manipulating the expression of *HY1* and *HY2* genes within different cells and tissues of the plant. Finally, it will be of  
15 particular interest to address the relationship of phytochrome chromophore biosynthesis and chlorophyll biosynthesis, not only because they share common biosynthetic intermediates but to determine how each pathway influences the other.

### **Methods.**

#### **Plant Materials**

20 *Arabidopsis thaliana* ecotypes Columbia (Col), Landsberg *erecta* (Ler), and Wassilewskija (Ws) were obtained from our laboratory stocks. Mutant strains used in this work were obtained from Maarten Koornneef for *hy2-1* (distributed as CS68 by the Arabidopsis Biological Stock Center, Columbus, OH; in *Ler* ecotype); from Jason Reed for *hy2-101* (EMS89S73S-E isolated originally by J. Reed; in Col ecotype), *hy2-102* (EMS195  
25 isolated by J. Reed; in Col ecotype), *hy2-103* (IAA<sup>R</sup>-7 isolated by Allison Wilson; in Col ecotype), *hy2-104* (IAA<sup>R</sup>-12 isolated by A. Wilson; in Col ecotype), *hy2-105* (γ10-9 isolated by J. Reed; in Col ecotype), and *hy2-106* (FN16-3 isolated by Aron Silverstone; in *Ler* ecotype); and from Nam-Hai Chua for *hy2-107* (segregated *hy2* from T-DNA lines in his laboratory; in WS ecotype). Plants were grown under long day conditions at 22°C in a  
30 growth chamber.

#### **Map-Based Cloning**

The *hy2-1* mutant was outcrossed with wild-type Col ecotype, and the mapping population was selected from F2 families with the long hypocotyl phenotype. Genomic DNA was prepared using a protocol described by Edwards *et al.* (1991) *Nucleic*

5 *Acids Res.* 19: 1349. We used cleaved amplified polymorphic sequence (CAPS) markers between Col and Ler (Konieczny and Ausubel (1993) *Plant J.* 4: 403-410), two CAPS markers (C6 and manganese-superoxide dismutase) in the Arabidopsis database, and seven new CAPS markers developed during this study. Primer sequences for polymerase chain reaction (PCR) amplification are listed here with the enzymes used for digestion indicated

10 in parentheses: c4523, 5'-ACA GCG AGA TTC AAA GGT CCA TTA ACC GGA-3' (SEQ ID NO:1) and 5'-GGG CTT ACA GTG ATA TCT GCA AGA CTT CTA -3' (HpaII) (SEQ ID NO:2); cMLP3E1, 5'-TAA TGC TTG CGA CAA ACA GG-3' (SEQ ID NO:3) and 5'-GTT CAT CTC AGG GCC AAA AA-3' (RsaI) (SEQ ID NO:4); cMXK7, 5'-GCT TTC AGA AAT CAG ACC TCA A-3' (SEQ ID NO:5) and 5'-CTG GTG TGG TTG ATC GAA

15 TCT-3' (DdeI) (SEQ ID NO:6); cMZB10, 5'-CTG CCA AGC TTC ATT TGG TT-3' (SEQ ID NO:7) and 5'-GCA GGA GCT GCA GAC AAT CT-3' (BsrI) (SEQ ID NO:8); cMZB10.18 (=HY2), 5'-CAA TGC AGG TTT AAC TTC AGC A-3' (SEQ ID NO:9) and 5'-CCA TGG GAA AGT CTG CAA AT-3' (DdeI) (SEQ ID NO:10); cF3L24, 5'-TCA AGC CCT TTT CCA ACA TC-3' (SEQ ID NO:11) and 5'-TTC CCC ATC TGA ACT CAA CC-

20 3' (HinfI) (SEQ ID NO:12); and cF8A24, 5'-AAT GAT GCA TGG TGT TGG TG-3' (SEQ ID NO:13) and 5'-GCT CGA GGA AAA GTC ATC CA-3' (MboI) (SEQ ID NO:14).

#### **Sequence Analysis of the HY2 Locus**

A pair of primers (5'-CGT TTG TCT CAC TGA AAC TG-3' (SEQ ID NO:15) and 5'-CAA TCA TCT TGA AAT GCA GA-3' (SEQ ID NO:16)) was used to

25 amplify 1.98-kb fragments of the MZB10.18 region from mutants and their corresponding wild-type plants. The PCR products were subjected directly to a cycle-sequencing protocol with several primers, and reactions were analyzed on an ABI373S sequencing apparatus (Applied Biosystems, Foster City, CA).

#### **Isolation of HY2 cDNA**

30 A cDNA library was constructed by K. Ando (Nara Institute of Science and Technology) from Col seedlings in  $\lambda$ ZAPII (Stratagene) according to the manufacturer's instructions. The DNA fragment containing MZB10.18 described above was used as a probe to screen ~300,000 cDNA clones by plaque hybridization. Several cDNA plasmids were recovered by in vivo excision according to the manufacturer's instructions.

### **RNA Isolation and Analysis**

RNA was isolated from 1-week-old whole Arabidopsis seedlings by the acid guanidinium thiocyanate-phenol-chloroform extraction method using Isogen (Nippon Gene, Tokyo, Japan). Total RNA (10 µg/lane) was electrophoresed on a 1.2% formaldehyde/agarose gel and transferred to a nylon membrane (Hybond-N; Amersham Corp.). Prehybridization and hybridization were then performed in Church hybridization solution (Church and Gilbert (1984)*Proc. Natl. Acad. Sci., USA*, 81: 1991-1995) using radioactive probes ( $3 \times 10^6$  to  $5 \times 10^6$  cpm/mL). A fragment of cDNA produced by EcoRI and XhoI digestion was used as a hybridization probe. Filters were washed under highly stringent conditions three times with  $1 \times$  SSC ( $1 \times$  SSC is 0.15 M NaCl and 0.015 M sodium citrate), 0.1% SDS at room temperature and twice with  $0.2 \times$  SSC, 0.1% SDS at 65°C for 15 min. To show equal loading of RNA, an rRNA probe was used for hybridization.

### **Subcellular Localization Experiment with Green Fluorescent Protein Fusion**

The coding region of *HY2* for the putative transit peptide and flanking amino acid residues (amino acids 1 to 62) isolated by PCR was cloned into pTH2XA, a modified green fluorescent protein (GFP) vector derived from 35SΩ-sGFP-S65T (Chiu *et al.* (1996) *Curr. Biol.* 6: 325-330). In pTH2XA, five glycine residues were included at the fusion junction to GFP (M. Takemura, unpublished data). The construct, which can express the *HY2* transit peptide fused to the N terminus of a modified GFP gene under the control of the cauliflower mosaic virus 35S promoter, was introduced into onion bulbs and tobacco leaves. The conditions of bombardment were the same as those described by Muramoto *et al.* (1999) *Plant Cell* 11: 335-347. Transient expression was observed after overnight incubation using confocal laser scanning microscopy (LSM510; Carl Zeiss, Jena, Germany).

### **Construction of the pGEX-mHY2 Expression Vector**

*mHY2*, the mature *HY2* gene without the predicted chloroplast transit peptide, was subcloned into the *Escherichia coli* expression vector pGEX-6-P1 (Amersham Pharmacia Biotech, Piscataway, NJ) to produce pGEX-mHY2. *mHY2* was amplified using the primers mHY2BglII fwd. (5'-GAA GAT CTG TCT CTG CTG TGT CGT ATA AGG-3', SEQ ID NO:17) and HY2SmaI rev. (5'-TCC CCC GGG TTA GCC GAT AAA TTG TCC TGT TAA ATC-3', SEQ ID NO:18), which contained BglII and SmaI sites (underlined),

5 respectively, and was cloned into BamHI-SmaI-digested pGEX-6-P1 to give pGEX-mHY2. The integrity of the construct was verified by restriction analysis and complete DNA sequencing of the insert (Davis Sequencing, Inc., Davis, CA). The constructed vector contains the *mHY2* sequence placed 3' to the glutathione-*S*-transferase (GST) gene of *Schistosoma japonicum* under the control of a *Ptac* promoter. A recognition sequence for  
10 PreScission protease, which is also a GST fusion protein, is located upstream of mHY2. Proteolytic cleavage yields the native Arabidopsis mHY2 with the five-amino acid N-terminal extension GPLGS.

### **Expression and Purification of Recombinant mHY2**

The Escherichia strain DH5 $\alpha$  containing pGEX-mHY2 was grown at 37°C  
15 in 500-mL batches of Luria-Bertani medium containing ampicillin (100  $\mu$ g/mL) to an OD<sub>578</sub> of 0.6. Cultures were induced by the addition of 1 mM isopropylthio- $\beta$ -galactoside and incubated for an additional 3 hr, and bacteria were harvested subsequently by centrifugation. The bacterial pellet from 3 liters of culture was resuspended in 20 mL of lysis buffer (50 mM Tris-HCl, pH 8.0, 100 mM NaCl, 0.05% Triton X-100, 1 mM DTT, 2  
20 mM benzamidine, 2 mM PMSF, leupeptin [2.0  $\mu$ g/mL], and pepstatin A [3  $\mu$ g/mL]) and disrupted with a French press (3  $\times$  20,000 p.s.i.). Cell debris were removed by centrifugation for 30 min at 100,000g. The resulting supernatant was loaded directly onto a glutathione-agarose (Sigma) column (1 cm  $\times$  3 cm) that had been equilibrated with 5 column volumes of PBS. Unbound protein was removed by washing the column with 5  
25 column volumes of PBS. GST-mHY2 fusion protein was eluted with 50 mM Tris-HCl, pH 8.0, containing 10 mM reduced glutathione. GST-mHY2-containing fractions were pooled and dialyzed overnight against cleavage buffer (50 mM Tris-HCl, pH 7.0, 100 mM NaCl, 1 mM EDTA, and 1 mM DTT). Digestion of the fusion protein was performed by adding 2 units of PreScission protease (Amersham Pharmacia Biotech) per 100  $\mu$ g of fusion protein  
30 and incubating at 4°C for 5 h. Removal of uncleaved fusion protein and excised GST tag was achieved by loading the digestion mixture onto a second glutathione-agarose column (1 cm  $\times$  3 cm). Recombinant mHy2 was detected in the flow through, analyzed by SDS-PAGE, and concentrated using Centriprep-10 concentrator devices (Amicon, Beverly, MA). One liter of bacterial culture yielded approximately 1 mg of purified protein.

### Determination of Protein Concentration

Protein concentration was determined using the method of Bradford (1976) *Anal. Biochem.* 72: 248-254, or by absorption at 280 nm for purified mHY2, where 1 absorption unit represents 0.64 mg/mL mHy2 (Gill and von Hippel (1989) *Anal. Biochem.* 182: 319-326).

### PΦB Synthase Activity Assay

All enzymes used for PΦB synthase assay were obtained from Sigma. For a 1-mL assay of PΦB synthase, the protein fraction to be assayed was diluted into 50 mM Tes-KOH, pH 7.3, containing an NADPH-regenerating system (6.5 mM glucose-6-phosphate, 0.82 mM NADP<sup>+</sup>, 1.1 units/mL glucose-6-phosphate dehydrogenase type XII from *Torula* yeast [EC 1.1.1.49]), a ferredoxin-reducing system (4.6 μM spinach ferredoxin, 0.025 units/mL spinach ferredoxin:NADP<sup>+</sup> oxidoreductase [EC 1.18.1.2]), and 10 μM BSA (fraction V, heat shock). Glucose-6-phosphate and NADP<sup>+</sup> were prepared as 100- and 25-mM stocks, respectively, in water; both were stored at 4°C. The glucose-6-phosphate stock was filter sterilized before storage. Glucose-6-phosphate dehydrogenase was prepared as a 500-unit/mL stock in 5 mM sodium citrate, pH 7.4, and stored at 4°C. Spinach ferredoxin:NADP<sup>+</sup> oxidoreductase was prepared as a 10-unit/mL stock with sterile water and stored at 4°C. BSA was made up as a 100-μM stock solution in 0.1 M potassium phosphate buffer, pH 7.4, and stored at either 4 or -20°C. The reaction was initiated by the addition of 5 μM (final concentration) purified biliverdin IXα (McDonagh and Palma (1980) *Biochem. J.* 189: 193-208) in 5 μL of DMSO. Assay mixtures were incubated in a 28°C water bath under green safe light or under subdued light for the desired amount of time. The assays were stopped by placing them on ice. Product analysis used a direct HPLC assay or a coupled assay after the addition of recombinant cyanobacterial apophytochrome 1 (Cph1) and difference spectroscopy (see below).

### Direct HPLC Assay

For the quantitative analysis of PΦB synthase activity, assay mixtures (outlined above) were loaded onto a Waters (Milford, MA) C<sub>18</sub> Sep-Pak Light (catalog No. WAT023501) preconditioned as follows: 3-mL wash with acetonitrile to wet the Sep-Pak,

5 3-mL wash with MilliQ water, and 3-mL wash with 50 mM 4-methylmorpholine/glacial  
acetic acid (pH 7.7). After the sample was loaded onto the Sep-Pak, it was washed with 3  
mL of 4-methylmorpholine/glacial acetic acid (pH 7.7) followed by 3 mL of 0.1% (v/v)  
trifluoroacetic acid. The Sep-Pak was then eluted with 2 mL of 100% acetonitrile. The  
eluate was dried using a Speed-Vac lyophilizer. The dried samples were analyzed by  
10 HPLC. Samples were first dissolved in 10  $\mu$ L of DMSO and then diluted with 200  $\mu$ L of  
the HPLC mobile phase (acetone:20 mM formic acid [50:50, v/v]). After the samples were  
dissolved, they were centrifuged briefly, passed through a 0.45- $\mu$ m polytetrafluoroethylene  
syringe filter, and chromatographed using a Varian (Palo Alto, CA) 5000 liquid  
15 through absorbance detector. Peak areas were quantitated using a 3365 Chemstation II  
(Hewlett-Packard, Waldbronn, Germany). The HPLC column used for all of the analyses  
was a Phenomenex (Torrance, CA) Ultracarb 5- $\mu$ m ODS (20) 4.6-mm  $\times$  250-mm analytical  
column with a 4.6-mm  $\times$  30-mm guard column of the same material. The mobile phase  
used with this column was acetone:20 mM formic acid (50:50, v/v). The flow rate was 0.8  
20 mL/min.

### **Coupled Difference Spectral Assay**

An alternative to the direct analysis of P $\Phi$ B synthase activity was the  
coupled, or indirect, assay. This assay was based on the method outlined previously (Terry  
and Lagarias (1991) *J. Biol. Chem.* 266: 22215-22221). The assay described above for  
25 P $\Phi$ B synthase was performed as before, but instead of working up the sample by Sep-Pak,  
an aliquot of recombinant apophytochrome (Cph1 from *Synechocystis* sp. PCC 6803) was  
added to the sample. The sample was incubated for an additional 20 to 30 min at room  
temperature under green safe light, and then a difference spectrum was taken. The method  
for difference spectroscopy was described previously (Terry and Lagarias (1991) *J. Biol.*  
30 *Chem.* 266: 22215-22221).

### **Example 2**

#### **Functional Genomic Analysis of the HY2 Family of Ferredoxin-Dependent Bilin Reductases from Oxygenic Photosynthetic Organisms**

Phytobilins are linear tetrapyrrole precursors of the light-harvesting  
35 prosthetic groups of the phytochrome photoreceptors of plants and the phycobiliprotein

5 photosynthetic antennae of cyanobacteria, red algae, and cryptomonads. Previous biochemical studies have established that phytobilins are synthesized from heme via the intermediacy of biliverdin IX $\alpha$  (BV), which is reduced subsequently by ferredoxin-dependent bilin reductases with different double-bond specificities. By exploiting the sequence of phytochromobilin synthase (HY2) of Arabidopsis, an enzyme that catalyzes the  
10 ferredoxin-dependent conversion of BV to the phytochrome chromophore precursor phytochromobilin, genes encoding putative bilin reductases were identified in the genomes of various cyanobacteria, oxyphotobacteria, and plants. Phylogenetic analyses resolved four classes of HY2-related genes, one of which encodes red chlorophyll catabolite reductases, which are bilin reductases involved in chlorophyll catabolism in plants. To test the catalytic  
15 activities of these putative enzymes, representative HY2-related genes from each class were amplified by the polymerase chain reaction and expressed in *Escherichia coli*. Using a coupled apophytochrome assembly assay and HPLC analysis, we examined the ability of the recombinant proteins to catalyze the ferredoxin-dependent reduction of BV to phytobilins. These investigations defined three new classes of bilin reductases with distinct  
20 substrate/product specificities that are involved in the biosynthesis of the phycobiliprotein chromophore precursors phycoerythrobilin and phycocyanobilin. Implications of these results are discussed with regard to the pathways of phytobin biosynthesis and their evolution.

### **Introduction.**

25 Phytobilins are linear tetrapyrrole molecules synthesized by plants, algae, and cyanobacteria that function as the direct precursors of the chromophores of the light-harvesting phycobiliproteins and of the photoreceptor phytochrome (Beale (1993) *Chem. Rev.* 93: 785–802; Hughes and Lamparter (1999) *Plant Physiol.* 121: 1059–1068). The pathways of phytobin biosynthesis have been elucidated by biochemical fractionation of  
30 plant and algal extracts, by overcoming a blocked step with exogenous putative intermediates, and by analysis of linear tetrapyrrole-deficient mutants (Beale and Cornejo (1991) *J. Biol. Chem.* 266: 22328–22332; Beale and Cornejo (1991) *J. Biol. Chem.* 266: 22333–22340; Beale and Cornejo (1991) *J. Biol. Chem.* 266: 22341–22345 Terry *et al.* (1993) *Arch. Biochem. Biophys.* 306: 1–15). These studies indicate that the biosynthesis of  
35 phytobilins shares common intermediates with heme and chlorophyll biosynthetic pathways

5 to the level of protoporphyrin IX, at which point the latter two pathways diverge by metalation with iron or magnesium (Beale (1993) *Chem. Rev.* 93: 785–802).

Phytobilins are derived from heme, which is converted to biliverdin IX (BV), the first committed intermediate in their biosynthesis. In red algae, cyanobacteria, and plants, this interconversion is accomplished by ferredoxin-dependent heme oxygenases that are related in sequence to the mammalian heme oxygenase (Cornejo *et al.* (1998) *Plant J.* 15: 99–107; Davis *et al.* (1999) *Proc. Natl. Acad. Sci., USA*, 96: 6541–6546; Muramoto *et al.* (1999) *Plant Cell* 11: 335–347). Although they catalyze the same reaction, mammalian heme oxygenases use an NADPH-dependent cytochrome P450 reductase to generate reducing power for heme catabolism (Maines (1988) *FASEB J.* 2: 2557–2568).

15 The metabolic fate of BV differs in mammals, cyanobacteria, and plants, with BV being metabolized by different reductases with unique double-bond specificities (Fig 1). Mammalian biliverdin IX reductase (BVR), an NAD(P)H-dependent enzyme that catalyzes the two-electron reduction of BV at the C10 methine bridge to produce bilirubin IX (BR), was the first of these enzymes to be discovered (Maines and Trakshel (1993) *Arch. Biochem. Biophys.* 300: 320–326). A similar enzyme, encoded by the gene *bvdR*, was identified recently in cyanobacteria (Schluchter and Glazer (1997) *J. Biol. Chem.* 272: 13562–13569). Cyanobacteria and red algae also possess novel ferredoxin-dependent bilin reductases for the synthesis of the linear tetrapyrrole precursors of their phycobiliprotein light-harvesting antennae complexes (Beale and Cornejo (1991) *J. Biol. Chem.* 266: 22328–22332; Beale and Cornejo (1991) *J. Biol. Chem.* 266: 22333–22340; Beale and Cornejo (1991) *J. Biol. Chem.* 266: 22341–22345; Cornejo *et al.* (1998) *Plant J.* 15: 99–107).

Primarily on the basis of studies with the red alga *Cyanidium caldarium*, these investigators proposed that the biosynthesis of the two major phycobiliprotein chromophore precursors, phycoerythrobilin (PEB) and phycocyanobilin (PCB), requires two ferredoxin-dependent bilin reductases and several double-bond isomerases. The first bilin reductase catalyzes the two-electron reduction of BV at the C15 methine bridge to produce the BR isomer 15,16-dihydrobiliverdin (DHBV), whereas the second bilin reductase catalyzes the conversion of 15,16-DHBV to 3Z-PEB, a formal two-electron reduction of the C2 and C31 diene system. In *C. caldarium*, an additional enzyme mediates the isomerization of 3Z-PEB to 3Z-PCB, both of which appear to be isomerized to their corresponding 3E isomers before assembly with the nascent phycobiliprotein apoproteins



5 (Beale and Cornejo (1991) *J. Biol. Chem.* 266: 22328–22332; Beale and Cornejo (1991) *J. Biol. Chem.* 266: 22333–22340; Beale and Cornejo (1991) *J. Biol. Chem.* 266: 22341–22345).

More recent studies lend support for a similar pathway of PCB and PEB synthesis in cyanobacteria (Cornejo and Beale (1997) *Photosynth. Res.* 51: 223–230). In contrast with mammals and phycobiliprotein-containing organisms, plants and green algae reduce BV to 3Z-PΦB by the ferredoxin-dependent enzyme PΦB synthase, which targets the 2,3,31,32-diene system for reduction (Terry *et al.* (1995) *J. Biol. Chem.* 270: 11111–11118; Wu *et al.* (1997) *J. Biol. Chem.* 272: 25700–25705). In plants, 3Z-PΦB is isomerized to its 3E isomer, which appears to be the immediate precursor of the phytochrome chromophore (Terry *et al.* (1995) *J. Biol. Chem.* 270: 11111–11118). The green alga *Mesotaenium caldarium* possesses a second bilin reductase activity that catalyzes the reduction of the 18-vinyl group of PΦB to produce 3Z-PCB (Wu *et al.* 1997 R15R15). These investigations also revealed that 3E-PCB is the natural phytochrome chromophore precursor in this organism.

20 Despite the extensive biochemical analysis of the phytobilin biosynthetic pathways in plants, algae, and cyanobacteria, the low levels of bilin reductase expression have hindered efforts to clone these enzymes. Using a genetic approach the HY2 locus of *Arabidopsis*, which encodes the enzyme PΦB synthase was cloned (Example 1).

The studies reported here were undertaken to identify HY2-related genes in the protein and nucleic acid databases. Using cloning, expression, and biochemical characterization, our investigations revealed three new classes of ferredoxin-dependent bilin reductases with either unique substrate or product specificities.

## **Results.**

### **The HY2-related gene family in cyanobacteria, oxyphotobacteria, and plants**

30 Example 1 describes the cloning of the HY2 gene of *Arabidopsis*. Using the deduced protein sequence of HY2, TBLASTN, BLASTP, and PSI-BLAST searches (Altschul *et al.* (1990) *J. Mol. Biol.* 215: 403–410; Altschul *et al.* (1997) *Nucleic Acids Res.* 25: 3389–3402) were performed to identify putative bilin reductases in the nonredundant National Center for Biotechnology Information database, in CyanoBase (Nakamura *et al.* 2000 R20R20), and in the Joint Genome Institute Microbial Genome database

- 5 (http://spider.jgi-psf.org/JGI\_microbial/html). These searches identified 15 putative proteins from various photosynthetic bacteria and two known proteins from plants.

Figure 10 shows a multiple sequence alignment of this family of proteins using CLUSTAL W (Higgins *et al.* (1996) *Meth. Enzymol.* 266: 383–402), hand adjustment with MEME (Bailey and Elkan (1995) pp. 21–29 In: *Proc. Third Internat. Conf. on*  
 10 *Intelligent Systems for Molecular Biology*, Menlo Park, CA: American Association of Artificial Intelligence Press), and highlighting with GENEDOC (http://www.psc.edu/biomed/genedoc). This alignment revealed regions of strong similarity interspersed with highly diverged regions, with an average pairwise similarity score of 25%. No sequence similarity of these proteins was observed with mammalian biliverdin  
 15 reductases.

On the basis of the biochemical data presented here, we name these *HY2*-related cyanobacterial loci after their roles in the biosynthesis of PCB (*i.e.*, *pcyA*) and PEB (*i.e.*, *pebA* and *pebB*). One of these proteins, the product of locus *slr0116* (*i.e.*, *pcyA*) in the genome of the cyanobacterium *Synechocystis sp* PCC6803, appears to be part of an operon  
 20 with a putative response regulator located 62 bp upstream (Ashby and Mullineaux (1999) *FEMS Microbiol. Lett.* 181: 253–260). Interestingly, this response regulator belongs to the OmpR subfamily for which a mutation (*ycf27*) was shown to cause a reduced energy transfer from the phycobilisomes to photosystem I (Ashby and Mullineaux (1999) *FEMS Microbiol. Lett.* 181: 253–260). *pcyA*-related open reading frames (orfs) also were found in  
 25 the oxyphotobacterium *Prochlorococcus sp.* MED4 (CCMP1378), which is also known as *Prochlorococcus marinus* MED4, in the marine cyanobacterium *Synechococcus sp* WH8102, and in the nitrogen-fixing, heterocyst-forming filamentous cyanobacteria *Anabaena sp* PCC7120 and *Nostoc punctiforme*. Among the other identified *HY2*-related genes are two orfs, *orf236* and *orf257*, from the marine cyanobacterium *Synechococcus sp*  
 30 WH8020 that lie adjacent to each other within the major phycobiliprotein gene cluster (Wilbanks and Glazer (1993) *J. Biol. Chem.* 268: 1226–1235; Wilbanks and Glazer (1993b) *J. Biol. Chem.* 268: 1236–1241). These orfs, which encode the proteins Ycp2\_SYNPY and Ycp3\_SYNPY, appear to be part of a three-gene operon containing an upstream orf of unknown function, *orf200*. A similar operon was identified in *Synechococcus sp* WH8102.  
 35 The genomes of *N. punctiforme* and *Prochlorococcus*, both the MED4 and SS120 (CCMP1375) subspecies, also contain similar operons. In contrast to the *N. punctiforme*

5 and *Anabaena* operons, an upstream orf in the *Prochlorococcus* operons exhibits a striking similarity to the ferredoxin-dependent heme oxygenase gene HY1 (Davis *et al.* (1999) *Proc. Natl. Acad. Sci., USA*, 96: 6541–6546; Muramoto *et al.* (1999) *Plant Cell* 11: 335–347) and its homologs in the cyanobacterium *Synechocystis* sp PCC6803 (Cornejo *et al.* (1998) *Plant J.* 15: 99–107). On the basis of their roles in PEB biosynthesis shown in this study, we  
10 name these ORFs *pebA* and *pebB*.

PSI-BLAST iterations also identified a weak relatedness of HY2 to the red chlorophyll catabolite reductase (RCCR) from barley and *Arabidopsis*. RCCR is involved in chlorophyll catabolism and catalyzes the ferredoxin-dependent reduction of the linear tetrapyrrole, red chlorophyll catabolite (RCC), to yield the primary fluorescent chlorophyll  
15 catabolite (Wüthrich *et al.* (2000) *Plant J.* 21: 189–198). These investigators showed that RCCR was incapable of reducing BV to either bilirubin or PB (Wüthrich *et al.* (2000) *Plant J.* 21: 189–198). Interestingly, the sequence similarity between RCCR and the other HY2-related proteins is so weak that TBLASTN searches using the two RCCR sequences failed to identify HY2 or other HY2-related proteins present in the publicly available databases  
20 (Wüthrich *et al.* (2000) *Plant J.* 21: 189–198). This divergence undoubtedly reflects the unusual substrate specificity of the RCCR for bilins derived from chlorophyll catabolism.

Phylogenetic analysis of the HY2-related family of proteins was performed using a heuristic parsimony search with a modified PAM250 weighting matrix and the program PAUP\* version 4.0 (see Methods). A single tree obtained with this analysis (Fig  
25 11) revealed four clades of HY2-related proteins with strong bootstrap support: PcyA, PebA, PebB, and RCCR. We noted that *HY2* lies within in the PebB clade.

### **Recombinant HY2-Related Proteins Are Bilin Reductases**

The HY2-related cyanobacterial orfs were amplified by polymerase chain reaction and cloned into the *Escherichia coli* expression vector pGEX-6P-1, which is  
30 similar to the vector described for mHY2 (Example 1). With this vector, the proteins were expressed as glutathione S-transferase (GST) fusions, which enabled their purification by affinity chromatography. The GST tag was removed via site-specific protease digestion, which resulted in an additional five to eight N-terminal amino acids due to the cloning strategy. Figure 12 shows SDS-PAGE results of purified recombinant protein  
35 representatives of the PcyA, PebA, PebB, and HY2 subfamilies. One liter of bacterial

5 culture yielded between 1 and 10 mg of soluble recombinant protein depending on which protein was expressed. The deduced molecular masses of the recombinant processed proteins, confirmed by SDS-PAGE (Fig 12), are as follows: *Anabaena* PcyA, 28.7 kD; *Synechocystis* PcyA, 28.9 kD; *Synechococcus* sp WH8020 PebA, 28 kD; *Synechococcus* sp WH8020 PebB, 30.3 kD; and *Arabidopsis* mHY2, 33.4 kD.

10 To determine whether the recombinant HY2-related proteins possess bilin reductase activity, we used a coupled holophytochrome assembly assay to analyze crude protein extracts from *E. coli* expressing these proteins for their ability to convert BV to phytobilins under standard PB synthase assay conditions. Figure 13A shows that crude bacterial lysates containing GST fusions of mHY2 and PcyA\_SYNY3 (PcyA\_ANASP; not  
15 shown) all exhibited BV reductase activities, yielding phytobilin products that could combine with the cyanobacterial phytochrome Cph1 apoprotein (apoCph1) to yield phytochrome difference spectra. The bilin metabolites incubated with apoCph1 resulted in different maxima and minima, suggesting that the various proteins reduced BV to distinct products. Both PcyA-containing extracts produced a BV metabolite(s) that gave spectra  
20 identical to those of the PCB adduct of apoCph1, with difference maxima at 655 nm and minima at 705 nm (Yeh *et al.* (1997) *Science* 277: 1505–1508). Figure 13A shows that both difference peaks of the mHY2 metabolites were markedly red shifted, with maxima at 670 and 730 nm, which is characteristic of the PΦB adduct of apoCph1 (Yeh *et al.* (1997) *Science* 277: 1505–1508; Example 1). Identical results were obtained using the purified  
25 recombinant HY2 and PcyA proteins (data not shown). Similarly, *E. coli* extracts lacking HY2 or PcyA proteins failed to metabolize BV to bilin products that could functionally assemble with apoCph1 (data not shown).

In contrast to the results for PcyA and HY2, no phytochrome difference spectrum was observed when the BV metabolites from reactions containing PebA\_SYNPY,  
30 PebB\_SYNPY, or a 1:1 mixture of the two *Synechococcus*-derived proteins were incubated with apoCph1. To determine whether fusion to GST is responsible for inhibiting the enzyme activity of these proteins, GST was removed by protease digestion and the full-length proteins were purified (Figure 12). Neither the purified proteins nor the 1:1 mixture of PebA and PebB were able to convert BV to a bilin product(s) that yielded a photoactive  
35 adduct with apoCph1 (Fig 13A). The observation that coincubation of a 1:1 ratio of PebA and PebB with BV elicited a color change of the assay mixture from bluish-green to pink

5 suggested that these proteins converted BV to bilins unable to form a photoconvertible holophytochrome. It is noteworthy that this pronounced color change was not observed when either PebA or PebB was assayed separately. This strongly implied that the PebA/PebB mixture could convert BV to PEB, the precursor of the phycobiliprotein C-phycoerythrin.

10 To test this hypothesis, BV-derived bilin metabolites from PebA, PebB, and PebA/PebB were incubated with apoCph1, and the mixtures were analyzed spectrofluorometrically for the production of the fluorescent PEB-apoCph1 "phytofluor" adducts (Murphy and Lagarias (1997) *Curr. Biol.* 7: 870–876). Only the PebA/PebB product mixture yielded a highly fluorescent compound, whose excitation and emission  
15 spectra were consistent with the formation of a phytofluor (Figure 13B). This result suggested that PebA and PebB were both required for the conversion of BV to PEB.

#### **HPLC Reveals Distinct Substrate/Product Specificity for Each Member of the HY2 Family**

HPLC analysis was performed to identify the bilin metabolites of the HY2  
20 family members using a chromatographic system that is able to separate 3E and 3Z isomers of PB, PCB, and PEB. As shown in Example 1, recombinant mHY2 efficiently reduced BV by two electrons to yield a mixture of both isomers of PB. In comparison, both PcyA proteins converted BV to a mixture of the 3E and 3Z isomers of PCB, a four-electron reduction (Fig 14). A time-course experiment was performed and revealed no evidence for  
25 other colored bilin intermediates (data not shown). Incubation of PebA\_SYNPY with BV resulted in the formation of an early eluting product that was detectable only at 560 nm and not at 380 nm (Fig 14). Optical spectroscopy revealed that this product had an absorption maximum at 575 nm in acetone:20 mM formic acid (50:50, v/v) (data not shown). Based on its absorption spectrum (data not shown), early retention time, and results shown below,  
30 this product was determined to be 15,16-DHBV. A similar absorption spectrum for DHBV has previously been published (Beale and Cornejo (1991b) *J. Biol. Chem.* 266: 22333–22340). In contrast to PebA\_SYNPY, PebB\_SYNPY was unable to metabolize BV (Figure 14). Identical results were observed with the *N. punctiforme* PebA and PebB homologs (data not shown).

5           A mixture of PebA and PebB effectively converted all of the BV to two colored pigments, one purple (retention time 9 min) and the other pink (retention time 10.5 min), whose retention times differed from that of 15,16-DHBV (Figure 14). Both bilin metabolites have absorption maxima in acetone:20 mM formic acid (50:50, v/v) near 580 nm (data not shown). Because PebB could not metabolize BV, these results suggest either  
10   that the 15,16-DHBV product of PebA was metabolized by PebB to the purple and pink bilins or that PebB forms a complex with PebA to alter its product profile. That 15,16-DHBV was a substrate for PebB was demonstrated by incubation of PebB with HPLC-purified 15,16-DHBV. In this case, the same two bilin products were observed (data not shown). HPLC coelution experiments showed the purple and pink pigments to be the 3E  
15   and 3Z isomers of PEB, respectively (data not shown). Both pigments are chemically stable in the HPLC mobile phase, eluting as single peaks after purification and reinjection. Moreover, both HPLC-purified pigments form phytofluors upon incubation with apoCph1, indicating that these are configurational isomers of PEB. HPLC-purified 15,16-DHBV from the PebA-mediated reduction of BV, however, was unable to form a fluorescent adduct with  
20   apoCph1 (data not shown).

          Biochemical studies of ferredoxin-dependent bilin reductases from algae and plants indicated that the 3Z isomers of PEB, PCB, and PB were the primary metabolites of these enzymes, with the formation of the 3E isomer requiring distinct bilin isomerase(s) (Beale and Cornejo (1991a) *J. Biol. Chem.* 266: 22328–22332; Beale and Cornejo (1991b)  
25   *J. Biol. Chem.* 266: 22333–22340; Beale and Cornejo (1991c) *J. Biol. Chem.* 266: 22341–22345; Cornejo and Beale (1997) *Photosynth. Res.* 51: 223–230). Our results show that both bilin isomers are produced with recombinant HY2, PcyA, and PebA/PebB proteins. We believe that the production of the 3E isomers occurred because of the presence of glutathione in the assay mixture and because of heating in the Speed-Vac concentrator. In  
30   this regard, glutathione-mediated 3Z to 3E isomerization of phycobilins has been reported for bilin reductases from *C. caldarium* (Beale and Cornejo (1991) *J. Biol. Chem.* 266: 22341–22345). Preliminary experiments performed with GST fusion proteins that did not come in contact with reduced glutathione or with proteins that were eluted from the affinity column by protease digestion greatly increased the relative amount of 3Z isomers produced  
35   (data not shown). Heating that occurred during concentration also contributed to the formation of the 3E isomers. If the drying time was reduced, only 3Z isomers were detected

5 (data not shown). Therefore, we conclude that in all cases the 3Z isomers are the primary reaction product of these reductases and that the production of the 3E isomers occurs by non-enzyme-mediated side reactions caused by heat and reduced glutathione.

### **The HY2 Family of Bilin Reductases Are Ferredoxin Dependent**

10 All of the reductive interconversions of BV and 15,16-DHBV presented here were dependent on reduced ferredoxin, which necessitated the inclusion of ferredoxin:NADP<sup>+</sup> oxidoreductase and an NADPH-regenerating system in the assay mixture. Indeed, none of the reduced bilin metabolites were detectable via HPLC when either ferredoxin or the NADPH-regenerating system was omitted from the assay mixture (data not shown). These results are in agreement with the ferredoxin dependence of the  
15 bilin reductases from plants and algae (Beale (1993) *Chem. Rev.* 93: 785–802). Thus, this family of proteins constitutes a new class of bilin:ferredoxin oxidoreductases (EC 1.3.7.n).

### **Discussion.**

Using a combination of protein-based pattern searches of genomic databases, phylogenetic analysis, and biochemical characterization, these investigations establish that  
20 the HY2 family of ferredoxin-dependent bilin reductases can be subdivided into five classes: PcyA, PebA, PebB, HY2, and RCCR families (Figure 11). This classification system is supported by the distinct substrate preference and double-bond regiospecificity of representative members of each bilin reductase subfamily. PcyA, PebA, and HY2 all recognize BV as a substrate, yet each yields different bilin products. PebB and RCCR  
25 possess unique bilin substrates (*i.e.*, 15,16-DHBV and RCC, respectively), and neither metabolizes BV (Wüthrich *et al.* (2000) *Plant J.* 21: 189–198; this study). Biochemical analyses of representatives of the three new classes of bilin reductases identified here, PcyA, PebA, and PebB, document their involvement in the biosynthesis of the phycobiliprotein chromophore precursors PCB and PEB.

### **The PcyA Family of Ferredoxin-Dependent BV Reductases Plays a Key Role in PCB Biosynthesis**

30 In this investigation, we have documented that the *pcyA* genes of the cyanobacteria *Synechocystis sp* PCC6803, *Anabaena sp* PCC7120, and *N. punctiforme* (data not shown) encode bilin reductases that catalyze the four-electron reduction of BV to 3Z-

5 PCB. PCB is the precursor of the chromophores of the phycobiliproteins phycocyanin and allophycocyanin, which are abundant in all three cyanobacteria. PcyA enzymes are atypical bilin reductases because all others catalyze two-electron reductions. Formally, these enzymes catalyze two-electron reductions of both the A and D rings of BV; however, we have not detected the production of semi-reduced intermediates such as PB and 18<sup>1</sup>,18<sup>2</sup>-  
10 DHBV. Thus, it appears that the partially reduced intermediates are tightly bound to the enzyme. The direct conversion of BV to PCB in these cyanobacteria is in contrast to the proposed pathways of PCB biosynthesis in the red alga *C. caldarium*, which involves the intermediacy of PEB (Beale (1993) *Chem. Rev.* 93: 785–802), and in the green alga *M. caldarium*, in which 3Z-PΦB is an isolable intermediate (Wu *et al.* (1997) *J. Biol. Chem.*  
15 272: 25700–25705). pcyA-related genes also are present in the oxyphotobacterium *Prochlorococcus* sp MED4, an unanticipated observation in view of the lack of phycobiliproteins in this organism. Phylogenetic analyses place this oxyphotobacterial protein in the PcyA clade of PCB:ferredoxin oxidoreductases. We were also able to clone the *Prochlorococcus* sp MED4 pcyA gene and express it as an N-terminal GST fusion. We  
20 determined that recombinant PcyA\_PROME was able to reduce BV to PCB in our standard phytochrome-based assay (data not shown). It therefore possesses the same enzymatic activity as all other studied PcyA enzymes.

### **peb Operons Encode Bilin Reductases Involved in PEB Biosynthesis**

We have observed that the *pebA* and *pebB* genes of the cyanobacteria  
25 *Synechococcus* sp WH8020 and *N. punctiforme* encode bilin reductases that catalyze the conversions of BV to 15,16-DHBV and 15,16-DHBV to 3Z-PEB, respectively (Fig 1). *PebA* therefore is a 15,16-DHBV:ferredoxin oxidoreductase, whereas *PebB* is a 3Z-PEB:ferredoxin oxidoreductase. Both activities are consistent with the pathway of PEB biosynthesis in the red alga *C. caldarium* (Beale (1993) *Chem. Rev.* 93: 785–802). The two  
30 *peb* genes also are found in the same operon in both phycoerythrin-producing cyanobacteria, and their close association with the major phycobiliprotein gene clusters supports their role in phycobilin biosynthesis (Wilbanks and Glazer (1993) *J. Biol. Chem.* 268: 1236–1241). We hypothesize that *PebA* and *PebB* function as a dual enzyme complex, in view of the synergistic metabolism of BV observed when the two enzymes are  
35 coincubated. A *peb* operon is not present in the genome of the cyanobacterium



- 5 *Synechocystis* sp PCC6803, an organism that lacks phycoerythrin. This strongly suggests that PCB is synthesized in this cyanobacterium via the PcyA-dependent pathway, as opposed to the PEB pathway found in *C. caldarium* (Beale (1993) *Chem. Rev.* 93: 785–802). In this regard, biochemical analyses of crude extracts from *Synechocystis* sp PCC6803 provide no evidence for the production of PEB (Cornejo and Beale (1997) *Photosynth. Res.* 51: 223–230).

The MED4 and SS120 subspecies of the oxyphotobacteria *Prochlorococcus* also possess *peb* operons very similar to those of *Synechococcus* sp WH8020 and WH8102, except that the former possess upstream genes related to heme oxygenase. This strongly suggests that both oxyphotobacterial subspecies can synthesize PEB. In this regard, genes encoding the  $\alpha$  and  $\beta$  subunits of a novel phycoerythrin have been identified in the SS120 subspecies of *Prochlorococcus* (Hess et al. 1996 R29R29, Hess et al. 1999 R30R30). It also has been shown that this unusual phycoerythrin plays a role in light harvesting in this ecotype (Lokstein et al. (1999) *Biochim. Biophys. Acta* 1410, 97–98), which is adapted for photoautotrophic growth at great ocean depths where light is limited. This observation is consistent with the lack of phycoerythrin genes in the high light-adapted MED4 ecotype. Although the enzymatic activities of *Prochlorococcus* PebA and PebB have not been determined experimentally, our phylogenetic reconstructions suggest that these proteins may be functional orthologs of the *Synechococcus* and *Nostoc* enzymes. Further analysis of the bilin biosynthetic pathways in *Prochlorococcus* and marine cyanobacteria such as *Synechococcus* sp WH8020 will be interesting, because the shorter wavelength-absorbing phycourobilin chromophores are major constituents of their phycoerythrins (Ong and Glazer (1991) *J. Biol. Chem.* 266: 9515–9527; Hess et al. (1996) *Proc. Natl. Acad. Sci., USA*, 93: 11126–11130). Although we have identified PCB and PEB biosynthetic enzymes in both organisms, it remains to be determined whether either these or other enzymes play a role in phycourobilin biosynthesis.

#### **Phycobilin Isomerases: Are They Necessary?**

PcyA, HY2, and PebB mediate bilin reductions that yield the 3Z isomer of their respective products. Because numerous studies have established that the more thermodynamically stable 3E isomers are substrates for assembly of the phycobiliprotein and phytochrome holoproteins, it has been proposed that there are unique 3Z/3E isomerases

5 that mediate this interconversion (Beale (1993) *Chem. Rev.* 93: 785–802; Terry *et al.* (1993) *Arch. Biochem. Biophys.* 306: 1–15). It should be noted that the 3Z isomer of PB has been shown to be a substrate for apophytochrome (Terry *et al.* (1995) *J. Biol. Chem.* 270: 11111–11118); however, these investigators suggested that isomerization to the 3E isomer is necessary to yield the correct stereochemistry of the holophytochrome chromophore. Such  
10 an isomerase activity has been identified in extracts of the red alga *C. caldarium*; however, this reaction also can be mediated by reduced glutathione (Beale and Cornejo (1991) *J. Biol. Chem.* 266: 22333–22340). For this reason, the need for a 3Z/3E isomerase has been questioned. All of the hy mutant loci have now been cloned from *Arabidopsis*, and none of these genes appear to encode a bilin isomerase. Thus, the isomerization of 3Z-PΦB may  
15 occur chemically or may be catalyzed by a genetically redundant family of bilin isomerases.

X-ray crystallographic analyses of phycobiliproteins have revealed that the stereochemistries of the thioether linkages to distinct cysteine residues are not all the same (Schirmer *et al.* (1987) *J. Mol. Biol.* 196: 677–695; Schmidt *et al.* (1987) *Z. Naturforsch.* 42C, 845–848). Therefore, we propose that the different stereochemistries arise from the  
20 use of the 3Z and 3E isomers of the phycobilin precursor as substrates for assembly to distinct cysteinyl moieties. Beale and Cornejo (1991) *J. Biol. Chem.* 266: 22333–22340, have identified a bilin isomerase that catalyzes the conversion of 3Z-PEB to 3Z-PCB in *C. caldarium*, which appears to be the sole pathway for PCB biosynthesis in this organism. More recently, a lyase/isomerase from the cyanobacterium *Mastigocladus laminosus* was  
25 described that is involved in both the isomerization of PCB to phycoviolobilin and its covalent attachment to apophytoerythrocyanin (Zhao *et al.* (2000) *FEBS Lett.* 469: 9–13). On the basis of these results and the diversity of bilin isomers found in phycobiliproteins from marine cyanobacteria, cryptomonads, and oxyphotobacteria (Ong and Glazer (1991) *J. Biol. Chem.* 266: 9515–9527; Hess *et al.* (1996) *Proc. Natl. Acad. Sci., USA*, 93: 11126–  
30 11130; Wedemayer *et al.* (1996) *Photosynth. Res.* 48: 163–170), it is likely that numerous bilin isomerases are present in these oxygen-evolving photosynthetic organisms.

### **Molecular Evolution of the HY2 Family of Bilin Reductases**

A single phylogenetic tree that is well supported with bootstrap replicates was obtained for the HY2 family (Fig 11). This tree delineates four clades of bilin  
35 reductases, which is in good agreement with the enzymes' double-bond specificity for

5 reduction. HY2 appears most closely related to the PebB clade of enzymes that catalyze a  
reduction of 15,16-DHBV to PEB. We predict that phytochromobilin synthases, because of  
their exquisite BV substrate specificity, will form a distinct clade when HY2 orthologs from  
other plant species are identified. The relatedness of HY2 and PebA enzymes is reasonable  
because both families mediate reduction of the vinyl pyrrole A ring to form the ethylidene  
10 moiety. We speculate that these two classes arose from a common ancestor that used BV as  
a substrate. This notion is based on the observation that the PebA family of bilin reductases  
also recognizes BV as a substrate.

Unlike HY2 and PebB, members of the PebA family target the 15,16 double  
bond of BV for reduction. To evolve the PebA and PebB/HY2 subfamilies, we envisage a  
15 duplication of a ferredoxin-dependent BV reductase gene and subsequent divergence in a  
marine cyanobacterium growing in a light-limited environment. Such an environment  
would provide the selection pressure favoring evolution of the biosynthetic pathway for  
PEB, whose incorporation into phycoerythrin extends the light-harvesting wavelength range  
of their phycobilisomes. Depending on the rooting of the HY2 family tree, the comparative  
20 branch lengths of the PebA and PebB/HY2 families on the phylogenetic tree suggest that  
the A ring reductases are more ancient, with the 15,16 reductases evolving more recently.  
On the basis of these inferences, we speculate that a cyanobacterial progenitor of plant  
chloroplasts possessed a bilin reductase with an A ring reductase regiospecificity. The  
progenitor of present day cyanobacteria likely would have possessed the ability to  
25 synthesize PCB, an essential component of their allophycocyanin-containing phycobilisome  
core. Thus, the common *pebA/pebB* ancestor might have resembled present-day *pcyA*  
genes, which encode atypical BV reductases that catalyze the four-electron reduction of BV  
to PCB. To date, *pcyA* genes appear to be present in all cyanobacteria, whereas a *peb*  
operon is lacking in the phycoerythrin-deficient cyanobacterium *Synechocystis* sp  
30 PCC6803.

The role of the *pebA*, *pebB*, and *pcyA* genes in *Prochlorococcus* sp MED4  
remains a mystery. Members of this genus are distinguished by the presence of integral  
membrane antennae complexes that contain divinyl chlorophyll *a*<sub>2</sub> and *b*<sub>2</sub> and by the lack of  
phycobilisomes (Partensky *et al.* (1999) *Microbiol. Mol. Biol. Rev.* 63: 106–127).  
35 Functional phycoerythrins have been detected only for the SS120 subspecies. As such,  
these organisms have been thought by some to be descendants of the class of prokaryotic

5 photosynthetic organisms whose endosymbiosis led to higher plant chloroplasts. Phylogenetic analyses using 16S rRNA indicate that this probably is not the case, because *Prochlorococcus* species appear more similar to marine *Synechococcus* species than to chloroplasts (Urbach *et al.* (1998) *J. Mol. Evol.* 46: 188–201). These analyses also suggest that *Prochlorococcus* evolved more recently from a phycobilisome-containing ancestor that  
10 resembled a marine *Synechococcus* species. The need for *pebA*, *pebB*, and *pcyA* genes for phycobilin biosynthesis in this ancestor is self-evident, and such genes may not yet have been lost from *Prochlorococcus* species. It is conceivable that these BV reductases are required to make bilin chromophore precursors of light receptors, such as the phytochromes (Hughes and Lamparter (1999) *Plant Physiol.* 121: 1059–1068). Although phytochrome-like genes are abundant in some cyanobacterial genomes, none are present in the genome of *Prochlorococcus* sp MED4 (data not shown). Alternatively, BV reductases may be needed to drive heme oxygenase, whose role in iron metabolism is well documented (Poss and Tonegawa (1997a) *Proc. Natl. Acad. Sci., USA*, 94: 10919–10924; Poss and Tonegawa (1997b) *Proc. Natl. Acad. Sci., USA*, 94: 10925–10930; Richaud and Zabulon (1997) *Proc.*  
20 *Natl. Acad. Sci., USA*, 94: 11736–11741; Schmitt (1997) *J. Bacteriol.* 179: 838–845).

In addition to the bilin reductases involved in phytobilin biosynthesis, a separate class exists of bilin reductases that are involved in chlorophyll degradation (Hörtensteiner *et al.* (1998) *J. Biol. Chem.* 273, 15335–15339; Wüthrich *et al.* (2000) *Plant J.* 21: 189–198). The pathway of chlorophyll degradation that occurs during plant  
25 senescence is similar to the heme degradation pathway (Matile and Hörtensteiner (1999) *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 50: 67–95). After dephytylation and magnesium removal, the chlorophyll macrocycle ring is opened by a monooxygenase that has yet to be cloned (Hörtensteiner *et al.* (1998) *J. Biol. Chem.* 273, 15335–15339). This is followed by a ferredoxin-dependent reduction of the bilin product catalyzed by the RCCR (Hörtensteiner  
30 *et al.* (2000) *Plant Biol.* 2: 63–67; Wüthrich *et al.* (2000) *Plant J.* 21: 189–198). RCCRs are the most diverged members of the ferredoxin-dependent bilin reductase family. Indeed, these enzymes have markedly different substrate specificities. It is notable that RCCRs catalyze a reduction very similar to that mediated by the PebA family (*i.e.*, a 15,16 double-bond reduction). The structural determinants that are responsible for RCCR's unique  
35 substrate specificity and double-bond regioselectivity will be interesting to discover. Presumably, chlorophyll catabolism would be important for chlorophyll-containing

5 prokaryotes; however, to date, RCCR genes are not readily identifiable in the genomes of any photosynthetic prokaryotes. It is possible that RCCR genes were lost, or alternately, that they evolved more recently from an HY2-like gene in the chloroplast endosymbiont progenitor, because they are found in cryptogams and plants (Hörtensteiner *et al.* (2000) Plant Biol. 2: 63–67).

### 10 Mechanistic Implications

Ferredoxin-dependent bilin reductase catalyzes two- and four-electron reductions of linear tetrapyrroles. Because ferredoxin is a one-electron carrier, these enzymes are mechanistically quite different from the NAD(P)H-dependent BVR/BvRD family of BV reductases. Preliminary analyses to date have failed to identify a metal or  
15 flavin cofactor in any of the recombinant enzymes reported here, suggesting that electrons are transferred directly to the bilin moiety, possibly via reduction of an amino acid residue within the enzyme. Although this finding suggests the presence of bilin radical intermediates, additional experiments are needed to assess this hypothesis. The oxygen sensitivity of RCCR supports the hypothesis that bilin radicals, which react with molecular  
20 oxygen, are produced during RCC catalysis (Wüthrich *et al.* (2000) *Plant J.* 21: 189–198).

Examination of highly conserved residues in the entire HY2 family and those within each of the five classes of bilin reductases provides information regarding residues important to the protein structure, ferredoxin interaction site, and substrate/product specificity. In this regard, only a small number of residues are conserved in the entire HY2  
25 family of enzymes. These include hydrophobic residues at positions 137, 157, 158, 256, and 314, Pro-151, Phe-221, Ser-222, and Asp-171 (Figure 10). The notable lack of conserved basic residues suggests that the propionyl moieties of the bilin substrates do not form salt linkages with the enzymes. The conserved hydrophobic residues proline and phenylalanine are likely to be involved in overall protein structure (*i.e.*, folding).

30 Alternately, they may form hydrophobic interactions with conserved regions of the various bilin substrates. The loss-of-function *hy2-1* and *hy2-104* alleles of phytochromobilin synthase from *Arabidopsis* support the critical role of Pro-151 in HY2's structure. The conserved serine and aspartate residues likely play catalytic roles, such as hydrogen bonding with the substrate and/or substrate protonation to make the bound bilin a better electron  
35 acceptor.

5               Despite the wide divergence of the HY2 family, we believe that these conserved residues indicate that the active sites of all members of this class are similar. We speculate that the distinct double-bond reduction specificities of the BV reductases (*i.e.*, PcyA, PebA, HY2), the 15,16-DHBV reductases (*i.e.*, PebB), and the RCCR families reflect the positioning of the respective substrates within the catalytic pocket. Because the A/B and  
10 C/D rings of BV are very similar but not identical, it is conceivable that the substrate binding sites of the PebA and HY2 enzymes are tailored to position BV in opposite orientations, favoring electron transfer to the bilin C/D ring or A ring, respectively. If this is true, then the PebB class might tether its 15,16-DHBV substrate in an orientation similar to that of the HY2 family, whereas RCC might be bound to RCCR in a manner similar to that  
15 in which BV is bound to PebA. Future studies will address the unique substrate/product specificity using domain swapping, site-directed mutagenesis, synthetic biliverdin substrates, and x-ray crystallography.

### **Biotechnological Implications**

              The availability of genes for bilin reductases that mediate the biosynthesis of  
20 PΦB, PCB, and PEB provides us with useful tools for numerous biotechnological applications. The ability to engineer the biosynthesis of PEB in any BV-producing organism is now feasible via the introduction of one or two genes. In this way, phytofluors potentially can be produced in any ferredoxin-containing organism. Coexpression of bilin reductase genes with apophytochromes should enable us to produce holophytochromes in  
25 bacteria and yeast. This will facilitate not only three-dimensional structural analysis of phytochrome but also the reconstruction of phytochrome signaling in a nonplant system in which we can exploit the power of molecular genetic analyses. This approach has proven invaluable for the structure-function analysis of the steroid hormone receptor family. By introducing the *pcyA* gene into wild-type and chromophore-deficient mutant plants, we also  
30 should be able to change the wavelength specificity of phytochrome, which may favorably alter plant growth and development in the field environment. Introduction of the *pebA* and *pebB* genes into plants potentially will shunt the conversion of BV to PEB, yielding photomorphogenetically challenged plants with fluorescent phytochromes. This would be especially useful for the analysis of the temporal and spatial patterns of phytochrome  
35 expression in plants.

## 5 Methods.

### Reagents

All chemicals, including glutathione agarose, were purchased from Sigma (St. Louis, MO) and were American Chemical Society grade or better. Restriction enzymes and Taq polymerase were from Gibco BRL (Cleveland, OH). HPLC-grade acetone and 80% formic acid were purchased from Fisher Scientific (Pittsburgh, PA). The expression vector pGEX-6P-1 and PreScission protease were obtained from Amersham Pharmacia Biotech (Piscataway, NJ). Centricon-10 concentrator devices were purchased from Amicon (Beverly, MA).

### Bioinformatics

15 Protein and nucleic acid database searches were performed using programs at publicly available World Wide Web sites. Preliminary sequence data were obtained from the Department of Energy Joint Genome Institute ([http://spider.jgi-psf.org/JGI\\_microbial/html/](http://spider.jgi-psf.org/JGI_microbial/html/)). Multiple sequence alignments were performed using the programs CLUSTAL W (Higgins et al. 1996 R21R21), GENEDOC (20 <http://www.psc.edu/biomed/genedoc>), and MEME (Bailey and Elkan (1995) pp. 21-29 In Proceedings of the Third Inter-national Conference on Intelligent Systems for Molecular Biology, Menlo Park, CA: American Association of Artificial Intelligence Press) to guide hand alignments. Phylogenetic analysis of the HY2-related family of proteins based on the alignment shown in Fig 10 was conducted using a heuristic parsimony search with a 25 modified PAM250 weighting matrix (Dayhoff *et al.* (1978) Pp 345-352 In: *Atlas of Protein Sequences and Structure*, M.O. Dayhoff, ed, Washington, DC: National Biomedical Research Foundation) using the program PAUP\* version 4.0 (Swofford (1993) *J. Gen. Physiol.* 102: 9A).

Because there are negative values in the PAM250 matrix, the most negative 30 penalty was set equal to zero, and all other values were increased correspondingly. Scores for transitions to and from gaps were not defined in the original matrix; they were set equal to the most costly transition (25) defined in the matrix. Characters 1 to 65 and 323 to 368 in the alignment were excluded from our analysis because they correspond to N- and C-terminal extensions not common to all members of the HY2 family (*i.e.*, plastid transit 35 peptide found on HY2 and red chlorophyll catabolite reductase (RCCR), C-terminal

- 5 extension found only on HY2). For *Hordeum vulgare* RCCR, missing characters 65 to 116 were replaced with question marks, which were weighted as zero. A rescaled consistency index was used for character weighting.

### **Construction of Expression Vectors**

- HY2-related genes from *Synechocystis sp* PCC6803, *Synechococcus sp*  
 10 WH8020, and *Anabaena sp* PCC7120 were amplified from chromosomal DNA via polymerase chain reaction using the following primers, which contained the indicated and underlined restriction sites: *Synechocystis* pcyA, BamHI fwd: 5'-AAG GAT CCA TGG CCG TCA CTG ATT TAA G-3' (SEQ ID NO:19), SalI rev: 5'-ACG CGT CGA CTA TTA TTG GAT AAC ATC AAA TAA GAC-3' (SEQ ID NO:20); *Synechococcus* pebA,  
 15 EcoRI fwd: 5'-GGA ATT CAT CTT TGA TTC ATT TCT CAA TG-3' (SEQ ID NO:21), NotI rev: 5'-ATA GTT AGC GGC CGC TCA TTT GTG AGA GGA GGA GGC-3' (SEQ ID NO:22); *Synechococcus* pebB, EcoRI fwd: 5'-GGA ATT CAT CAC AAA TCA AAG ATT CAA AAG C-3' (SEQ ID NO:23), NotI rev: 5'-ATA GTT AGC GGC CGC TTA TAG ATC AAA AAG CAC AGT GTG G-3' (SEQ ID NO:24); and *Anabaena* pcyA, EcoRI fwd: 5'-  
 20 GGA ATT CAT CTC ACT TAC TTC CAT TCC CTC-3' (SEQ ID NO:25), NotI rev: 5'-ATA GTT AGC GGC CGC TTA TTC TGG -GA GAT CAA ATA AC-3' (SEQ ID NO:26). The polymerase chain reaction products were then cut with the indicated enzymes and inserted into similarly restricted pGEX-6P-1. The integrity of the plasmid constructs was verified by complete DNA sequence determination of the insert (Davis Sequencing, Davis,  
 25 CA). All of the constructs place the HY2-related gene downstream of and in frame with the glutathione S-transferase (GST) gene of *Schistosoma japonicum* under the control of a Ptac promoter. A recognition sequence for PreScission protease is located upstream of the cloned gene. Proteolytic cleavage yields the native protein with a small N-terminal extension. In all cases, the original initiation methionine was changed to an isoleucine.

### **Expression and Purification**

- Expression and purification were performed according to instructions supplied by the manufacturer (Amersham Pharmacia Biotech) and as described in Example 1. Between 1 and 10 mg of purified protein was obtained per liter of bacterial culture.



5           **Protein Determination**

Protein concentration was determined by the Bradford method with BSA as a standard (Bradford 1976 R48R48) or by measuring the absorbance at 280 nm and using the calculated 280 nm for each individual protein (Gill and von Hippel (1989) *Anal. Biochem.* 182: 319-326).

10           **Standard Bilin Reductase Activity Assay**

Assays for bilin reductase activity were performed as described for PΦB synthase (*see* Example 1).

**Direct HPLC Analysis**

              Bilin reductase assay mixtures were loaded onto a Waters (Milford, MA)  
15 C18 Sep-Pak Light preconditioned as follows: 3-mL wash with acetonitrile to wet the Sep-Pak, 3-mL wash with MilliQ water, and 3-mL wash with 50 mM 4-methylmorpholine/glacial acetic acid, pH 7.7. After the sample was loaded onto the Sep-Pak, it was washed with 3 mL of 4-methylmorpholine/glacial acetic acid, pH 7.7, followed by 3 mL of 0.1% (v/v) trifluoroacetic acid. The bilin metabolites were then eluted from the  
20 Sep-Pak with 2 mL of 100% acetonitrile. The eluate was dried using a Speed-Vac lyophilizer (Savant Instruments Inc., Farmingdale, NY), and the dried samples were analyzed by HPLC. Samples were first dissolved in 10 µL of DMSO and then diluted with 200 µL of the HPLC mobile phase (50:50 v/v acetone:20 mM formic acid). After the samples were dissolved, they were centrifuged briefly to collect the sample, passed through  
25 a 0.45-µm polytetrafluoroethylene syringe filter, and chromatographed using a Varian (Palo Alto, CA) 5000 liquid chromatograph. The HPLC column used for all of the analyses was a Phenomenex (Torrance, CA) Ultracarb 5-µm ODS20 4.6 x 250-mm analytical column with a 4.6 x 30-mm guard column of the same material. The mobile phase used with this column was acetone:20 mM formic acid (50:50, v/v). The flow rate was 0.8  
30 mL/min. The eluate was monitored at 560 nm for the first 11.5 min and at 380 nm for the remaining time using a Varian UV100 flow-through absorbance detector. Peak areas were quantitated using a Hewlett-Packard (Palo Alto, CA) model 3365 Chemstation II.

### Coupled Spectrophotometric and Spectrofluorometric Analysis

5 An aliquot of 20 µg of crude recombinant Cph1 apoprotein (Yeh and Lagarias (1998) *Proc. Natl. Acad. Sci., USA*, 95: 13976–13981) was added to 1 mL of bilin reductase assay mixture under green safelight. Mixtures were incubated for 30 min at room temperature to permit phytyobilin binding. Phytochrome difference spectra were obtained as  
10 described previously (Terry and Lagarias (1991) *J. Biol. Chem.* 266: 22215–22221). A spectrofluorometric assay was used to detect the formation of intensely fluorescent phycoerythrobilin (PEB) adducts of Cph1 (Murphy and Lagarias (1997) *Curr. Biol.* 7: 870–876). Emission spectra were obtained with an excitation wavelength of 545 nm using an SLM Aminco Bowman AB2 spectrofluorometer (Spectronic Instruments Inc., Rochester,  
15 NY).

### Example 3

#### Production of Functional Phytochrome in Living Cells

In the higher plants two enzymes are committed to the biosynthesis of phytyochromobilin PΦB - the chromophore precursors of phytyochrome. These enzymes are  
20 heme oxygenase (encoded by *HY1* in *Arabidopsis thaliana* (Muramoto *et al.* (1999) *Plant Cell*, 11: 335–347)), which catalyzes the ferredoxin-dependent conversion of heme to biliverdin IXα (BV), and phytyochromobilin:ferredoxin oxidoreductase (PΦB synthase) in *Arabidopsis* (encoded by *HY2* in *Arabidopsis*), which catalyzes the ferredoxin-dependent conversion of BV to PΦB. A homolog of the *HY1* protein, *HO1* which is encoded by  
25 Cyanobase Locus SLL1184 of the cyanobacterium *Synechocystis* sp. PCC 6803 has been shown to be a functional ferredoxin-dependent heme oxygenase. Here we show that co-expression of the biosynthetic enzymes *HO1* and *HY2* together with the cyanobacterial phytyochrome Cph1 yields the production of photoreversible holophytyochrome in the bacterium *Escherichia coli*, with spectroscopic properties consistent with the formation of a  
30 phytyochromobilin-adduct.

This work involved the production of synthetic operon comprised of *HO1* from *Synechocystis* sp. PCC6803 and the mature *HY2* coding region (*mHY2*) from *Arabidopsis thaliana* that lacks the plastid targeting sequence. The cloning of *HO1* and *mHY2* open reading frames into the plasmid pPROLarA122 (Clontech Laboratories) places  
35 this operon under regulatory control of a dual *Ara/Lac* promoter. Upon introduction of this

5 plasmid into *E. coli* cells harboring the Cph1-expression plasmid, pBAD/Cph1(514), in which Cph1(N514) is under regulatory control of a *Ara* promoter, the production of photoactive holophytochrome *in vivo* was determined.

### Methods.

#### Plasmid Construction.

10 The synthetic operon consisting of *HO1* and *pcyA* coding regions was cloned in the expression vector pPROlarA122 to produce the plasmid pPROlarA122/HO1-RBS-SLR0116 (SEQ ID NO:27) as follows. The *HO1* gene from *Synechocystis* sp. PCC6803 was first PCR amplified with the sense primer Pho1-S1K, 5'-ATC GGT ACC ATG AGT GTC AAC TTA GCT TC-3' (SEQ ID NO:28) (containing a KpnI restriction site) and  
15 antisense primer Pho1-ArB, 5'-ATT GGA TCC TTT CTC CTC TTT AAC TAG CCT TCG GAG GTG GCG A-3' (SEQ ID NO:29) (containing a synthetic ribosome binding site upstream of a BamHI restriction site) using chromosomal DNA from *Synechocystis* sp. PCC6803 as a template. The reaction was carried out using a standard reaction mix, Taq polymerase, and a 30 cycle run with an annealing temperature of 50°C. The gene was then  
20 cloned into TA cloning plasmid, pCR2. 1 (Invitrogen), producing plasmid pCR2. 1/HO1-RBS (not shown).

The synthetic operon consisting of *HO1* from *Synechocystis* sp PCC6803 (Cornejo *et al.* (1998) *Plant J.* 15: 99-107) and mHY2 coding regions was produced by cloning mHY2 into the plasmid pCR2. 1/HO1-RBS to produce plasmid pCR2.1/HO1-RBS-  
25 HY2. Specifically, the mHY2 cDNA from Arabidopsis was PCR-amplified using plasmid DNA from the clone pGEX-mHY2 (Example 2), which contains the full length mHY2 cDNA minus the transit peptide in a GST-fusion vector, with sense primer mHY2-EcoRV: 5'-CGG ATA TCA TGT CCC CTAT ACT A-3' (SEQ ID NO:30) and the antisense primer, mHY2-NotI: 5'-GCG CGG CCG CTT AGC CGA TAA ATT GTC C-3' (SEQ ID NO:31)  
30 under standard conditions. The reaction was carried out using a standard reaction mix, Pfu polymerase, and a 35 cycle run with an annealing temperature of 55°C. The PCR product was restricted with EcoRV and NotI and then subcloned in to the plasmid pCR2. 1/HO1-RBS to produce the plasmid pCR2.1/HO1-RBS-HY2. Finally pCR2. 1/HO1-RBS-HY2 was restricted with KpnI/NotI and the resulting fragment was ligated with KpnI/NotI

- 5 restricted pPROLarA122 (Clontech Laboratories) resulting in production of pPROLarA122/HO1-RBS-HY2.

**E. coli strains, media, and transformation.**

- The plasmid pPROLarA122/HO1-RBS-HY2 was transformed into *E. coli* strain LMG194 (Invitrogen) competent cells containing the apophytochrome expression  
10 plasmid, pBAD/Cph1(514). Dual ampicillin and kanamycin selection using minimal RM media was performed to isolate transformants.

**Protein expression.**

- The *E. coli* strain LMG194 containing both plasmids pBAD/Cph1(514) and pPROLarA122/HO1-RBS-HY2 was grown overnight at 37°C in 3 ml RM media containing  
15 25 µg/ml kanamycin and 50 µg/ml ampicillin. A 1 ml aliquot of this culture was transferred to 100 ml of RM media and grown at 37°C to an OD<sub>600</sub> of approximately 0.5. 50 ml of this culture was then transferred to 450 ml LB media containing 25 µg/ml kanamycin and 50 µg/ml ampicillin. IPTG was added to a final concentration of 1 mM to induce expression of the synthetic operon. After incubation for 1 h at 30°C, arabinose was added to a final  
20 concentration of 0.002% to induce expression of apoCph1. The culture was grown at 30°C for 5 h, after which time cells were collected by centrifugation and resuspended in 10 ml lysis buffer (50 mM Tris-HCl, pH 8.0, 100 mM NaCl, 0.05% v/v NP40, 2 µg/ml leupeptin, 2 mM benzamidine, 2 mM PMSF, 1mM DTT, 3 µg/ml pepstatin A). A cell lysate was obtained by lysing the cells at 10,000 psi with a French Press. After insoluble material was  
25 removed by centrifugation, the crude homogenate was placed on ice at 4°C and examined for holophytochrome spectrophotometrically.

**Protein Purification:**

- The crude soluble fraction was run over a Talon (Clontech) metal affinity chromatography column (5 ml bed volume), washed with 20 ml extraction/wash buffer, and  
30 eluted with 2 bed volumes 1x elution buffer (EW buffer containing 200 mM imidazole). The resulting solution was dialyzed overnight against 2 liters of 10 mM HEPES pH 7.5, and then concentrated using an Amicon ultrafiltration cell.

5 **Results & Discussion.**

After 5 h induction at 30°C, cultures containing both pBAD/Cph1(514) and pPROlarA122/HO1-RBS-HY2 plasmids turned blue-green. As shown in the difference spectrum (Figure 15), co-expression of pBAD/Cph1(514) and pPROlarA122/HO1-RBS-HY2 yielded crude cell extracts containing photoactive holophytochrome. The spectrum of  
10 purified holoCph1(N514) from these cells reveals absorption maxima for the Pr form at 660 nm and for the Pfr form at 710 nm, consistent with the formation of a phytochromobilin (PΦB) adduct *in vivo*, as opposed to the blue-shifted phycocyanobilin (PCB) adduct formed in Cph1(N514)-expressing *E. coli* cells coexpressing the PCB operon, *i.e.* HO1 and PcyA (Figure 16) (Yeh, *et al.* (1997) *Science*, 277: 1505-1508).

15 It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference in their entirety for all  
20 purposes.

5

## CLAIMS

**What is claimed is:**

1. An isolated HY2 family bilin reductase comprising an amino acid consensus sequence as illustrated in Figure 5 or in Figure 10 and having bilin reductase activity.
- 10 2. The bilin reductase of claim 1, wherein said bilin reductase is not hvrccr or atrccr1.
3. The bilin reductase of claim 1, wherein said bilin reductase is not rccr\_horvu or rccr\_arath.
- 15 4. The bilin reductase of claim 1, wherein said bilin reductase is not ycp2\_synpy or ycp3\_synpy.
5. The bilin reductase of claim 1, wherein said bilin reductase comprises at least 50% sequence conservation as shown in Figure 10.
6. The bilin reductase of claim 1, wherein said bilin reductase comprises at least 70% sequence conservation as shown in Figure 10.
- 20 7. The bilin reductase of claim 1, wherein said bilin reductase comprises at least 90% sequence conservation as shown in Figure 10.
8. The bilin reductase of claim 1, wherein said bilin reductase comprises at least 80% sequence conservation as shown in Figure 5.
9. The bilin reductase of claim 1, wherein said bilin reductase comprises  
25 at least 100% sequence conservation as shown in Figure 5.
10. The bilin reductase of claim 1, wherein said bilin reductase is PebA.
11. The bilin reductase of claim 1, wherein said bilin reductase is PebB.
12. A ferredoxin-dependent bilin reductase comprising at least 15% sequence identity with an enzyme selected from the group consisting of HY2\_ARATH,

- 5 YCP2\_SYNPY, YHP2\_PROMA, YHP3\_PROMA, YCP3\_SYNPY, SLR0116,  
PcyA\_ANASP, PcyA\_NOSPU, PcyA\_SYNY3, PcyA\_SYN8.1, PcyA\_PROME,  
PebA\_SYNPY, PebA\_SYN8.1, PebA\_PROMA, PebA\_PROME, PewbB\_NOSPU,  
HY2\_ARATH, RCCR\_ARATH, and RCCR\_HORVU, and where, when aligned with HY2,  
comprises conserved hydrophobic residues at position 137, 157, 158, 256, and 314.
- 10 13. The bilin reductase of claim 12, wherein said bilin reductase, when  
aligned with HY2, comprises a residue selected from the group consisting of Pro-151, Phe-  
221, Ser222, and ASP-171.
14. The bilin reductase of claim 13, wherein said bilin reductase, when  
aligned with HY2, comprises Pro-151, Phe-221, Ser-222, and ASP-171.
- 15 15. The bilin reductase of claim 12, wherein said bilin reductase is not  
hvrccr or atrccr1.
16. The bilin reductase of claim 12, wherein said bilin reductase is not  
rccr\_horvu or rccr\_arath.
17. The bilin reductase of claim 12, wherein said bilin reductase is not  
20 ycp2\_synpy or ycp3\_synpy.
18. The bilin reductase of claim 12, wherein said bilin reductase is not  
HY2.
19. An isolated bilin reductase having bilin reductase activity and  
comprising an amino acid sequence of polypeptide selected from the group consisting of  
25 HY2, athy2, slr0116, c362\_anab, ycp2-synpy, ycp3\_synpy, PcyA\_ANASP, PcyA\_NOSPU,  
PcyA\_SYNY3, PcyA\_SYN81, PcyA\_PROME, PebA\_SYNPY, PebA\_SYN81,  
PebA\_PROMA, PebA\_PROME, PebA\_NOSPU, PebB\_SYNPY, PebB\_SYN81,  
PebB\_PROMA, PebB\_PROME, PebB\_NOSPU, HY2\_ARATH, RCCR\_ARATH, and  
RCCR\_HORVU, or conservative substitutions thereof.
- 30 20. The bilin reductase of claim 19, wherein said bilin reductase  
comprises an amino acid sequence of a polypeptide selected from the group consisting of  
athy2, slr0116, c362\_anab, ycp2-synpy, ycp3\_synpy, PcyA\_ANASP, PcyA\_NOSPU,

5 PcyA\_SYNY3, PcyA\_SYN81, PcyA\_PROME, PebA\_SYNPY, PebA\_SYN81,  
PebA\_PROMA, PebA\_PROME, PebA\_NOSPU, PebB\_SYNPY, PebB\_SYN81,  
PebB\_PROMA, PebB\_PROME, PebB\_NOSPU, HY2\_ARATH, RCCR\_ARATH, and  
RCCR\_HORVU.

21. A method of converting a biliverdin to a phytobilin, said method  
10 comprising contacting a bilin reductase of claim 1, with a biliverdin whereby said biliverdin  
is converted to a phytobilin.

22. The method of claim 19, wherein said bilin reductase is a  
cyanobacterial bilin reductase.

23. The method of claim 19, wherein said bilin reductase is an algal bilin  
15 reductase.

24. The method of claim 19, wherein said bilin reductase is a plant bilin  
reductase.

25. The method of claim 21, wherein said bilin reductase is  
recombinantly expressed.

20 26. The method of claim 21, wherein said contacting is *ex vivo*.

27. The method of claim 21, wherein said contacting is in a cell and said  
bilin reductase is a heterologous polypeptide.

28. The method of claim 21, further comprising contacting said  
phytochromobilin with a second bilin reductase to produce a phytochrome.

25 29. The method of claim 21, further comprising contacting said  
phytochromobilin with a second bilin reductase to produce a phytofluor.

30. The method of claim 29, wherein said second bilin reductase is PebB.

31. The method of claim 21, wherein said bilin reductase is ycp2-snpy.

32. The method of claim 29, wherein said bilin reductase is ycp3-snpy.



- 5                   33.    A nucleic acid comprising a nucleic acid encoding a bilin reductase  
of any one of claims 1 through 20.
34.    The nucleic acid of claim 33, wherein said nucleic acid is a vector.
35.    A cell comprising a heterologous nucleic acid comprising a nucleic  
acid encoding a bilin reductase of any one of claims 1 through 20.
- 10               36.    The cell of claim 35, wherein said cell is selected from the group  
consisting of an algal cell, a plant cell, a yeast cell, a bacterial cell, an insect cell, and a  
mammalian cell.
37.    A nucleic acid comprising a nucleic acid that specifically hybridizes  
with a nucleic acid of any one of claims 1 through 20 under stringent conditions and that  
15   encodes a polypeptide having bilin reductase activity, wherein said nucleic acid does not  
encode an hvrcr or an atrcr polypeptide.
38.    The nucleic acid of claim 37, wherein said nucleic acid is a vector.
39.    A method of detecting expression of a polypeptide, said method  
comprising:  
20                    providing a cell comprising a nucleic acid encoding an  
apophytochrome; and a nucleic acid encoding a bilin reductase that produces a phytobilin  
that assembles with said apophytochrome to produce a phytofluor;  
                      and detecting an optical signal produced by said phytofluor.
40.    A method of producing a photoactive holophytochrome, said method  
25   comprising:  
                      co-expressing in a cell:  
                          a heme oxygenase;  
                          an apophytochrome;  
                          and a ferredoxin-dependent bilin reductase;  
30   whereby said cell produces said photoactive holophytochrome and where one or more of  
said apophytochrome and said ferredoxin-dependent bilin reductase are expressed by  
heterologous nucleic acids.

- 5                   41.     The method of claim 40, wherein said cell is selected from the group consisting of an algal cell, a yeast cell, a bacterial cell, a plant cell, an insect cell, and a mammalian cell.
42.     The method of claim 40, wherein said ferredoxin-dependent bilin reductase is an HY2 family bilin reductase.
- 10                  43.     The method of claim 40, wherein said apophytochrome and said ferredoxin-dependent bilin reductase are both expressed by heterologous nucleic acids.
44.     The method of claim 40, wherein said heme oxygenase is expressed by a heterologous nucleic acid.
45.     The method of claim 40, wherein said photoactive holophytochrome  
15 is not a phytofluor.
46.     The method of claim 45, wherein said ferredoxin-dependent bilin reductase is an HY2 family member.
47.     The method of claim 45, wherein said ferredoxin-dependent bilin reductase is *HY2*.
- 20                  48.     The method of claim 45, wherein said ferredoxin-dependent bilin reductase is *pcyA*.
49.     The method of claim 40, wherein said photoactive holophytochrome is a phytofluor.
50.     The method of claim 49, wherein said apophytochrome is expressed  
25 as a fusion protein with a protein that is to be labeled with said phytofluor.
51.     The method of claim 49, wherein said method comprises expressing the ferredoxin-dependent bilin reductase *pebA* or *pebB*.
52.     The method of claim 51, wherein said method comprises expressing both ferredoxin-dependent bilin reductase *pebA* and *pebB*.

5 53. The method of claim 51, wherein said cell is a bacterial cell.

54. The method of claim 40, wherein said method further comprises recovering said photoactive holophytochrome from said cell.

55. A cell comprising:

a heme oxygenase;

10 an apophytochrome;

and a ferredoxin-dependent bilin reductase;

whereby said cell produces a photoactive holophytochrome and where one or more of said apophytochrome and said ferredoxin-dependent bilin reductase are expressed by heterologous nucleic acids.

15 56. The cell of claim 55, wherein said cell is selected from the group consisting of an algal cell, a yeast cell, a bacterial cell, a plant cell, an insect cell, and a mammalian cell.

57. The cell of claim 55, wherein said ferredoxin-dependent bilin reductase is an HY2 family bilin reductase.

20 58. The cell of claim 55, wherein said apophytochrome and said ferredoxin-dependent bilin reductase are both expressed by heterologous nucleic acids.

59. The cell of claim 55, wherein said heme oxygenase is an endogenous heme oxygenase.

25 60. The cell of claim 55, wherein said heme oxygenase is expressed by a heterologous nucleic acid.

61. The cell of claim 55, wherein said photoactive holophytochrome is not a phytofluor.

62. The cell of claim 61, wherein said ferredoxin-dependent bilin reductase is an HY2 family member.

- 5                   63.     The cell of claim 61, wherein said ferredoxin-dependent bilin  
reductase is *HY2*.
64.     The cell of claim 61, wherein said ferredoxin-dependent bilin  
reductase is *pcyA*.
- 10                  65.     The cell of claim 55, wherein said photoactive holophytochrome is a  
phytofluor.
66.     The cell of claim 65, wherein said apophytochrome is expressed as a  
fusion protein with a protein that is to be labeled with said phytofluor.
67.     The cell of claim 65, wherein said cell comprises expressing the  
ferredoxin-dependent bilin reductase *pebA* or *pebB*.
- 15                  68.     The cell of claim 67, wherein said cell comprises expressing both  
ferredoxin-dependent bilin reductase *pebA* and *pebB*.
69.     The cell of claim 67, wherein said cell is a bacterial cell.
70.     A recombinant nucleic acid comprising:  
                          a nucleic acid encoding a heme oxidoreductase; and  
20                        a nucleic acid encoding and a ferredoxin-dependent bilin reductase;  
where said nucleic acid expresses a functional heme oxidoreducuse and a functional bilin  
reductase.
71.     The nucleic acid of claim 70, wherein said heme oxidoreductase and  
said bilin reductase are under control of the same promoter.
- 25                  72.     The nucleic acid of claim 71, wherein said promoter is a constitutive  
promoter.
73.     The nucleic acid of claim 71, wherein said promoter is an inducible  
promoter.
74.     The nucleic acid of claim 71, wherein said promoter is a tissue-  
30     specific promoter.


5                    75.     The nucleic acid of claim 70, wherein said nucleic acid is present in a cell.


                  76.     The nucleic acid of claim 75, wherein said cell is selected from the group consisting of an algal cell, a bacterial cell, a plant cell, a yeast cell, a mammalian cell, and an insect cell.

10                  77.     The nucleic acid of claim 75, wherein said nucleic acid comprises a gene selected from the group consisting of *HO1*, *HY2*, *PcyA*, *PebA*, and *PebB*.

                  78.     The nucleic acid of claim 77, wherein said nucleic acid comprises an *HO1* coding region and a *pcyA* coding region.

                  79.     The nucleic acid of claim 78, wherein said nucleic acid further  
15     comprises a *pcyB*.

**Fig. 1**

**Fig. 2**




Fig. 3A




```

gaattccaccacgtcaacdgtagctgtgcattccacgtgggggatgtgggacctatagttag 60
aoccatgaactcggacggatgttgaaatcattgtcgttgccaattggcgtttgtctcactga 120
aactgtgaaaattttatctcttttatagataaAGAATCTTGCTTTTTCAGTTTTCAGTA 180
TGAAGAAGAAATTGAAGAGAGTGTCCGAGGAAGGAGACCTTTGCTTTCAGTTTGTGAGTCT 240
TGTTGTAAATGGCTTTATCAATGGAGTTTGGGTTTCAATTGGGTCATGCTTCAAGGCACC 300
  N A L S M E F G F S I G S C F K A P
AAACCCACCTGTTCTAATCTCTGCAAGGCCCTAATAAGATCAATTCACGTTGAGAAGGAG 360
  N P P V L I S A S P N K I N F T L R R R
AAAGAAAAGATTCTTACTTAGAGTCTCTGCTGTGTGTATAGGAATTCGCAGAGTCTGC 420
  hy2-106 *****
  R K R F L L R V S A V S Y K E F A E S A
TTTAGAAGAAACCAGGAAAAGGATCGTTCTTGAACTTCACATCTCCAGgtatatgcaat 480
  L E E T R K R I V L E P S H L Q
taoatttoggtagttagtgggaggatttatattctcattgttttctgtgtgaattttg 540
ggtaaatgattgttagttagttagttagttagttagttagttagttagttagttagttag 600
ttatataagtaaaagttcagattttgtttttctaatcagaactgtttcagGAAAAGTA 660
  E K Y
TAGTAGCATGACAGGACTAGATGGTAAGACCGAACTTCAAATGCTTGCTTTTAAATCTTTC 720
  S S M T G L D G K T E L Q M L A F K S S
AAAGATTAGACTCTTGAGGAGTATGGCAATAGAGAATGAGACAATGCAGgttttaacttca 780
  K I R L L R S M A I E N E T M Q
gongtaccactgattgttttagttagttagttagttagttagttagttagttagttagttag 840
  hy2-105 *****
tcttoggtagttagttagttagttagttagttagttagttagttagttagttagttagttag 900
  hy2-1,hy2-104 T
  V F D F A G F M E P E Y D T P I F
TCTGTGCTAACTTTTTCACATCTACCAACGTTAACAATAGTTGTATTgttaagtattattct 960
  C A N F F T S T N V N I V V L
agttatgtggagttatcaggtctgtattgtccaaactgatgttcaatattttactgtat 1020
gttctctcttagGGACCTTAATCCTTTGCATCAGTTGACTGACCAGACGGATTACCAAGA 1080
  D L N P L M Q L T D Q T D Y Q D
CAAGTATTATAACAAGATAATGTCCATATATCACAAATATGCTGAGgtgaccacaagaat 1140
  R Y Y N X I M S I Y E K Y A E
acacacaaattactcaattgcaagtaaacctaatgtgaggtgttaaatgactgatcttgag 1200
atattatttgagACTTTCCCATGGGGAGGGAATTGACTGGTGAATCCATAAAGTTTTC 1260
  hy2-101 A
  T F P W G G K L T G E S I K F F
TCGCCCTTTGTGATGTGGACTAGGTTTTCGTCTAGCAAGAAAAACATAAGGCTTTGTTC 1320
  S P L V M W T R F S S S K E K H K A L F
TCTGCGCTTCTAGAGTACTATCAGgtatatactcagcgcccaaaagctaaaggttttattg 1380
  S A F L E Y Y Q
gaacatttgactgngaatttatcatcttcttctacagGCATGCTTGAGATGACAATCC 1440
  hy2-107 a
  A W L E M T I Q
AAGTGAGGAGGAGATGGAACCATCTCATGTGAGAGCCAATTGTGAGCACAAACACAAGT 1500
  V R E E M E P S H V R A H C E A Q H K Y
ACCTGACATGGCGAGCACAAAAGgtgatttcatttcttttgtgttaatttgcattgtttga 1560
  hy2-103 A
  L T W R A Q K
acagacactgtatctgtattgtttacaaatggatattgatttgggtgtttgcagGATCCTGGA 1620
  hy2-102 a
  D P G
CATGGTCTTCTTAAAGATTAGTAGGTGAAGCAAAGGCAAGgtatataaaagatttgatcc 1680
  H G L L K R L V G E A K A K
cattagtggtcccatattatttaattagcttgtgaagatgttgaaatgatttgaacaaatc 1740
agGAGCTGCTAAGGSAATTTCCTGTTCATGGGCTGGATGAGTTAGGCACAAAAACATTC 1800
  E L L R D F L F N G V D E L G T K T F I
TTGATTACTTTCCAGAGTACCAACAGAGATGGAAGTGTAAAGCGATAAAGCAAGTATCA 1860
  D Y F P E Y Q T E D G T V S D K R S I I
TTGGGAGTCAATGAAACTCGTCCATGGGATTTAACAGGCAATTTATCGGCTAACAAAT 1920
  G K S Y E T R P W D L T G Q F I G
GATATATGTGAACAAAGTCAGATTTCAGAGTCATCAACACAAGAGGACGTGAACCTTAGGGA 1980
AGTAGGAATAAGAAAGAGCAGCATGAGGAGTCTCTCAGGTCTATCTGCATTTCAGATGA 2040
TTGTTTGAAGTACCATGCATTGTAGTTTTCACAGTGTAGCTCTCAGCCCTTCATCAAAT 2100
GAGAATCCTCGAGTATGATATGATTTTAAATGAAGATGTATTGCTCTTactaatcaaca 2160


```

Fig. 3B






**Fig. 4A and Fig. 4B**

**Fig. 5**



**Fig. 6A - 6D**

**Fig. 7****Fig. 8**

**Fig. 9**

10/15

```

PCYA_ANASP : 20 * 40 60 80
PCYA_NOSPU : MSLTSPSLREQQHPHTRQIADQAEVWQHQLDLSYPHMPAEL : 43
PCYA_SYNY3 : MSFTSMPSLREQQHPHTRQIADQAEVWQHQLDLSYPHMPAEL : 43
PCYA_SYNA1 : MAVTSLTSSSLMPTDMPITQQALAAASWQS-LPKKPYQPEDEL : 46
PCYA_PROMA : MQSPPESSSSIVAPLPSLAETRGAMGLPKHPLDASDF : 42
PCYA_SYNPY : INLLSKSITKTKLIDPITLILQKQVORSALNCLNCLNPPKL : 44
PCYA_SYNPY : MDSVLNMBHSDITK-RGGSPLPPEGL : 27
PCYA_SYNA1 : PDPTLEPQTGIA-RGGISVEVAGL : 27
PCYA_PROMA : MNKLMLQDLHNMKRIIS-HGGKPIRVENGH : 31
PCYA_PROMA : TESLKNPKTNIE--DLDGASGISK : 25
PCYA_NOSPU : MINSQSPLENVALFLINETCMIAITYFHARVNSKSCSAKPFLEHELELFORDELQSRVHPGL : 64
PCYA_SYNPY : MTNQRKXSTDPVNIENWQPFLEDAIARLEG-LNVEPYPRDRP : 44
PCYA_SYNA1 : MSIDLASSLDPVQIPGWRWQPFLEDAASAKIF-TNPSYPHATF : 45
PCYA_PROMA : MIKRDNSLSKIDLRDWNTPFFNDVEXDSV-FEIEEYPHSHDF : 44
PCYA_PROMA : MLIQNTIFYSQEWKAKFKFISQLDN-YHCVEHKASDF : 40
PCYA_NOSPU : MNSERSDVTINQPFLDYAIAYMRSRLDLEPYPTGF : 37
HY2_ARATH : MALSMBPGHSIGSCFKAPN-PPVLISASPNKINFTRRRKKRFLIRVSASVKEFAESAEETRRK-IVLEPSHPQEKY : 77
RCCR_ARATH : HAMIFCNWLYSSSPSYLSPLTSKPSRFSNLRRAQSQSMEDHDLRRKFMFPYVSPTRKQMNVDLMSTENRLOS-QOLLPCNPPDW : 96
RCCR_HORVU :

```

```

PCYA_ANASP : 100 120 140 160 180
PCYA_NOSPU : GYVEGRLEGK- LTFENRCYQTPQFKNMHELAKEVNMNDILHCHMGRGR-GOISAAEASVHL-ERTL : 126
PCYA_SYNY3 : GYVEGRLEGK- LTFENRCYQTPQFKNMHELAKEVNMNDILHCHMGRGR-GOISAAEASVHL-ERTL : 126
PCYA_SYNA1 : GYVEGRLEGK- LTFENRCYQTPQFKNMHELAKEVNMNDILHCHMGRGR-GOISAAEASVHL-ERTL : 126
PCYA_PROMA : SSIEGQLEGDD- LLIRNELCCRGVKNMHELAKEVNMNDILHCHMGRGR-GOISAAEASVHL-ERTL : 123
PCYA_SYNPY : SNIISNEEGKE- LYIENEFPKAGFKNMHELAKEVNMNDILHCHMGRGR-GOISAAEASVHL-ERTL : 127
PCYA_SYNA1 : EECRSKSS- SVIQSWLMDVPGFKNMHELAKEVNMNDILHCHMGRGR-GOISAAEASVHL-ERTL : 109
PCYA_PROMA : EHNQSQKGS- STIQSWLMDVPGFKNMHELAKEVNMNDILHCHMGRGR-GOISAAEASVHL-ERTL : 109
PCYA_SYNPY : SERFSHKQD- TVLXSWLMDVPGFKNMHELAKEVNMNDILHCHMGRGR-GOISAAEASVHL-ERTL : 109
PCYA_PROMA : EFKENHND- SKYITKNNIWSQYKNMHELAKEVNMNDILHCHMGRGR-GOISAAEASVHL-ERTL : 109
PCYA_SYNPY : EFKVSDRGR- NPATIRSWCYQSQELKNMHELAKEVNMNDILHCHMGRGR-GOISAAEASVHL-ERTL : 109
PCYA_NOSPU : LOREDTGSKK- SIPVTTATWACKTEFKNMHELAKEVNMNDILHCHMGRGR-GOISAAEASVHL-ERTL : 109
PCYA_SYNA1 : LQEGSTGSKK- PVPVTTATWACKTEFKNMHELAKEVNMNDILHCHMGRGR-GOISAAEASVHL-ERTL : 109
PCYA_PROMA : LKESITGSRN- PVHVTTLTWAAPFKIKNMHELAKEVNMNDILHCHMGRGR-GOISAAEASVHL-ERTL : 109
PCYA_SYNPY : SYKSSYSGSKS- KKNINFTWGAHQKINFAPACINSNYSVNFNFKTSNTPPLVDFSLP-TSHLMDVDFSLP-VNQF : 127
PCYA_NOSPU : ESNASVVGKGN- QREVTTSYAQTAKLIDRAAHVQCTMSQVNFNFKTSNTPPLVDFSLP-TSHLMDVDFSLP-VNQF : 124
HY2_ARATH : SSMTGLDG- KTRQMLAFKSSKIELRSMALN-ETQVDFPAGNPEPSTPTFCANFTST-NVNVMMDLNLHQLTDQDY : 159
RCCR_ARATH : PNFNPNGSABASHIRSGDKSSPIDFVIGSWHCKIPTTSNMLISISGLNSSTKANVWHEHQSS-SKSLNLDLBRKDLVIMPDY : 181
RCCR_HORVU : MDENLQSSLSHCKVPS-AIDHISLFINLNASITAPAFHINELQGS-PTSMVETLPRKDLALHPEY : 66

```

Fig. 10


11/15



PCYA\_ANASP : PESYNSA \* 200 \* 220 \* 240 \* 260 \*  
 PCYA\_NOSPU : PESYTTA \* 199 :  
 PCYA\_SYNS1 : PAAYQXS \* 199 :  
 PCYA\_SYNS1 : PSGLTAD \* 202 :  
 PCYA\_SYNS1 : PSGLTAD \* 198 :  
 PCYA\_SYNS1 : PSGLTAD \* 199 :  
 PCYA\_SYNS1 : PSGLTAD \* 186 :  
 PCYA\_SYNS1 : PSGLTAD \* 186 :  
 PCYA\_SYNS1 : PSGLTAD \* 190 :  
 PCYA\_SYNS1 : PSGLTAD \* 185 :  
 PCYA\_SYNS1 : PSGLTAD \* 224 :  
 PCYA\_SYNS1 : PSGLTAD \* 213 :  
 PCYA\_SYNS1 : PSGLTAD \* 214 :  
 PCYA\_SYNS1 : PSGLTAD \* 213 :  
 PCYA\_SYNS1 : PSGLTAD \* 209 :  
 PCYA\_SYNS1 : PSGLTAD \* 202 :  
 PCYA\_SYNS1 : PSGLTAD \* 237 :  
 PCYA\_SYNS1 : PSGLTAD \* 271 :  
 PCYA\_SYNS1 : PSGLTAD \* 155 :


PCYA\_ANASP : QOIGA \* 280 \* 300 \* 320 \* 340 \* 360 \*  
 PCYA\_NOSPU : QONIGA \* 245 :  
 PCYA\_SYNS1 : QONIGA \* 245 :  
 PCYA\_SYNS1 : QONIGA \* 247 :  
 PCYA\_SYNS1 : QONIGA \* 243 :  
 PCYA\_SYNS1 : QONIGA \* 236 :  
 PCYA\_SYNS1 : QONIGA \* 235 :  
 PCYA\_SYNS1 : QONIGA \* 241 :  
 PCYA\_SYNS1 : QONIGA \* 236 :  
 PCYA\_SYNS1 : QONIGA \* 280 :  
 PCYA\_SYNS1 : QONIGA \* 257 :  
 PCYA\_SYNS1 : QONIGA \* 262 :  
 PCYA\_SYNS1 : QONIGA \* 257 :  
 PCYA\_SYNS1 : QONIGA \* 257 :  
 PCYA\_SYNS1 : QONIGA \* 255 :  
 PCYA\_SYNS1 : QONIGA \* 329 :  
 PCYA\_SYNS1 : QONIGA \* 319 :  
 PCYA\_SYNS1 : QONIGA \* 205 :



Fig. 10 cont'd



**Fig. 11**

*Fig. 12**Fig. 13A and 13B*

**Fig. 14**

**Fig. 15****Fig. 16**

## SEQUENCE LISTING

SEQ ID NO:27.

DNA sequence of expression plasmid pPROLarA122/HO1-RBS-SLR0116

10

CAGAATTCATTAAAGAGGAGAAAGGTACCATGAGTGTCAACTTAGCTTCCCAGTTGCGGGAAGGGACGAAAAAA  
TCCCCTCCATGGCGGAGAACGTCGGCTTTGTCAAATGCTTCCTCAAGGGCGTTGTGAGAGAAAAATTCCTACCG  
TAAGCTGGTTGGCAATCTCTACTTTGTCTACAGTGCCATGGAAGAGGAAATGGCAAAATTTAAGGACCATCCCA  
TCTCAGCCACATTTACTTCCCCGAACCTCAACCGCAAACAAAGCCTAGAGCAAGACCTGCAATTCTATTACGGC  
15 TCCAACCTGGCGGCAAGAAGTGAAAAATTTCTGCCGCTGGCCAAGCCTATGTGGACCGAGTCCGGCAAGTGGCCGC  
TACGGCCCCCTGAATTGTTGGTGGGCCATTCTACACCCGTTACCTGGGGGATCTTTCCGGCGGTCAAATTTCTCA  
AGAAAAATTGCCCAAAATGCCATGAATCTCCACGATGGTGGCAGAGCTTTCTATGAATTTGCCGACATTGATGAC  
GAAAAGGCTTTTAAAAATACCTACCGTCAAGCTATGAATGATCTGCCCATTTGACCAAGCCACCGCCGAACGGAT  
TGTGGATGAAGCCAATGACGCCTTTGCCATGAACATGAAAAATGTTCAACGAACCTTGAAGGCAACCTGATCAAGG  
20 CGATCGGCATTATGGTGTTC AACAGCCTCACCCGTCGCCGCGAGTCAAGGCAGCACCGAAGTTGGCCTCGCCACC  
TCCGAAGGCTAGTTAAAGAGGAGAAAGGATCCATGGCCGTCACCTGATTTAAGTTTGACCAATTCTTCCCTGATG  
CCTACGTTGAACCCGATGATTC AACAGTTGGCCCTGGCGATCGCCGCTAGTTGGCAAAGTTTACCCCTCAAGCC  
CTATCAATTGCCGGAGGATTTGGGCTACGTAGAAGGCCGCTTGAAGGGGAAAAGTTAGTGATTGAAAATCGGT  
GCTACCAAACGCCCCAGTTTCGCAAAATGCATTTGGAGTTGGCCAAGGTGGGCAAAGGGTTGGATATTCTCCAC  
25 TGTGTAATGTTTCTTGAGCCTTTATACGGTCTACCTTTGTTTGGCTGTGACATTTGTGGCCGGCCCCGGTGGAGT  
AAGTGCGGCTATTGCGGATCTATCCCCCACCCAAAGCGATCGCCAATTGCCCGCAGCGTACCAAAAATCATTTGG  
CAGAGCTAGGCCAGCCAGAAATTTGAGCAACAACGGGAATTTGCCCCCTGGGGAGAAATATTTTCTGAATATTGT  
TTATTCATCCGTCCCAGCAATGTCACTGAAGAAGAAAGATTTGTACAAAGGGTAGTGGACTTTTTGCAAATTCAT  
TTGTACCAATCCATCGTTGCCGAACCCCTGTCTGAAGCTCAAACCTTTGGAGCACCGTCAGGGGCAAATTCATT  
30 ACTGCCAACACAACAGAAAAATGATAAAACCCGTCGGGTACTGGAAAAGCTTTTGGGGAAGCTTGGGCGGAA  
CGGTATATGAGCCAAGTCTTATTTGATGTTATCCAATAATCTAGAGGCATCAAATAAAACGAAAGGCTCAGTCG  
AAAGACTGGGCCTTTTCGTTTATCTGTTGTTTGTGCGGTGAACGCTCTCCTGAGTAGGACAAATCCGCCGCCCTA  
GACCTAGGGGATATATTCCGCTTCTCGCTCACTGACTCGCTACGCTCGGTGCTTCGACTGCGGCGAGCGGAAA  
TGGCTTACGAACGGGGCGGAGATTTCCCTGGAAGATGCCAGGAAGATACTTAACAGGGAAGTGAGAGGGCCGCGG  
35 CAAAGCCGTTTTCATAGGCTCCGCCCCCTGACAAGCATCACGAAATCTGACGCTCAAATCAGTGGTGGCGA  
AACCCGACAGGACTATAAAGATACCAGGCGTTTCCCCCTGGCGGCTCCCTCGTGCCTCTCCTGTTCTCTGCCTT  
TCGGTTTACCGGTGTCAATCCGCTGTTATGGCCGCGTTTGTCTCATTCACGCCTGACACTCAGTTCGGGTAG  
GCAGTTCGCTCCAAGCTGGACTGTATGCACGAACCCCCGTTCAAGTCCGACCGCTGCGCCTTATCCGGTAACCTA  
TCGTCTTGAGTCCAACCCGGAAGACATGCAAAAGCACCCTGGCAGCAGCCACTGGTAATTGATTTAGAGGAG  
40 TTAGTCTTGAAGTCATGCGCCGTTAAGGCTAAACTGAAAGGACAAGTTTTGGTGACTGCGCTCCTCCAAGCCA  
GTTACCTCGGTTCAAAGAGTTGGTAGCTCAGAGAACCCTCGAAAAACCGCCCTGCAAGGCGGTTTTTTTCGTTTT  
CAGAGCAAGAGATTACGCGCAGACCAAAACGATCTCAAGAAGATCATCTTATTAATCAGATAAAATATTACTAG  
ATTTCAAGTGAATTTATCTCTTCAAATGTAGCACCTGAAGTCAGCCCCATACGATATAAGTTGTTACTAGTGCT

5 TGGATTCTCACCAATAAAAAACGCCCCGGCGGCAACCGAGCGTTCTGAACAAATCCAGATGGAGTTCTGAGGTCA  
TTACTGGATCTATCAACAGGAGTCCAAGCGAGCTCTCGAACCCCGAGTCCCGCTCAGAAGAAGTCTGTCAGAA  
GGCGATAGAAGGCGATGCGCTGCGAATCGGGAGCGGCGATACCGTAAAGCACGAGGAAGCGGTGAGCCCATTCG  
CCGCCAAGCTCTTCAGCAATATCACGGGTAGCCAACGCTATGTCTTGATAGCGGTCCGCCACACCCAGCCGGCC  
ACAGTCGATGAATCCAGAAAAGCGGCCATTTTCCACCATGATATTCGGCAAGCAGGCATCGCCATGGGTACGA  
10 CGAGATCCTCGCCGTCGGGCATGCGCGCCTTGAGCCTGGCGAACAGTTCTGGCTGGCGCGAGCCCTGATGCTCT  
TCGTCCAGATCAFCCTGATCGACAAGACCGGCTTCCATCCGAGTACGTGCTCGCTCGATGCGATGTTTCTGCTTG  
GTGGTCGAATGGGCAGGTAGCCGGATCAAGCGTATGCAGCCGCCGATTCGATCAGCCATGATGGATACTTTCT  
CGGCAGGAGCAAGGTGAGATGACAGGAGATCCTGCCCCGGCACTTCGCCCAATAGCAGCCAGTCCCTTCCCGCT  
TCAGTGACAACGTCGAGCACAGCTGCGCAAGGAACGCCCGTCTGTGGCCAGCCACGATAGCCGCGCTGCCTCGTC  
15 CTGCAGTTCATTACAGGSCACCGGACAGGTGCGTCTTGACAAAAAGAACCAGGCGCCCCCTGCGCTGACAGCCGGA  
ACACGGCGGCATCAGAGCAGCCGATTTGTCGTGTTGTGCCCAGTCATAGCCGAATAGCCTCTCCACCCAGCGGCC  
GGAGAACCTGCGTGCAAATCCATCTTGTTCATCATGCGAAACGATCCTCATCCTGTCTCTTGATCAGATCTTGA  
TCCCTTGCGCCATCAGATCCTTGCGGGCAAGAAAGCCATCCAGTTTACTTTGCAGGGCTTCCCAACCTTACCAG  
AGGGCGCCCCAGCTGGCAATTCGACGTCGTGTGTGGAATTGTGAGCGGATAACAATTTACACAGGGCCCCCTCG  
20 ACACCGAGGAGAATGTCAAGAGGCGAACACACAACGCTCTTGAGCGCCAGAGGAGGAACGAGCTAAAACGGAGC  
TTTTTTGCCCTGCGTGACCAGATCCCGGAGTTGGAAAACAATGAAAAGGCCCCCAAGGTAGTTATCCTTAAAAA  
AGCCACAGCATACATCCTGTCCGTCCAAGCAGAGGAGCAAAAGCTCATTTCTGAAGAGGACTTGTGCGGAAAC  
GACGAGAACAGTTGAAACACAAACTTGAACAGCTACGGAACCTTGTGCGTAAGGAAAAGTAAGGAAAACGATT  
CCTTCTAACAGAAATGTCCTGAGCAATCACCTATGAACTGTCGACTCGAGCATAGCATTTTTATCCATAAGATT  
25 AGCGGATCTAACCTTTTACAAATTGTGAGCGCTCACAATTATGATAGATTCAATTGTGAGCGGATAACAATTTAC  
A

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
13 December 2001 (13.12.2001)

PCT

(10) International Publication Number  
**WO 01/094548 A3**

(51) International Patent Classification<sup>7</sup>: C12N 9/02,  
1/20, 15/00, C07H 21/04

Gregory, A.: 2516 Temple Drive, Davis, CA 95616 (US).  
**MONTGOMERY, Beronda, L.:** 728 W. 17th Street,  
Bloomington, IN 47404 (US).

(21) International Application Number: PCT/US01/18326

(22) International Filing Date: 5 June 2001 (05.06.2001)

(74) Agent: **BERLINER, Robert:** Fulbright & Jaworski  
L.L.P., 29th Floor, 865 South Figueroa Street, Los Ange-  
les, CA 90017-2571 (US).

(25) Filing Language: English

(81) Designated States (*national*): CA, JP.

(26) Publication Language: English

(84) Designated States (*regional*): European patent (AT, BE,  
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,  
NL, PT, SE, TR).

(30) Priority Data:

60/210,286	8 June 2000 (08.06.2000)	US
60/271,758	26 February 2001 (26.02.2001)	US
Not furnished	29 May 2001 (29.05.2001)	US

**Published:**

— with international search report  
— before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

(71) Applicant: **THE REGENTS OF THE UNIVERSITY  
OF CALIFORNIA** [US/US]; 1111 Franklin Street, 12th  
Floor, Oakland, CA 94607-5200 (US).

(88) Date of publication of the international search report:  
11 July 2002

(72) Inventors: **LAGARIAS, John, Clark:** 1853 Imperial  
Avenue, Davis, CA 95616 (US). **KOCHI, Takayuki:**  
8916-5 Takayama, Daigakusyukusya A-404, Ikoma,  
Nara 630-0101 (JP). **FRANKENBERG, Nicole:** 1224  
F. Street #2, Davis, CA 95616 (US). **GAMBETTA,**

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.



**WO 01/094548 A3**

(54) Title: HY2 FAMILY OF BILIN REDUCTASES

(57) **Abstract:** This invention identifies a novel family of bilin reductases. Designated herein HY bilin reductases, the enzymes of this invention are useful in a wide variety of contexts including but not limited to the conversion of biliverdins to phytobilins and the assembly of holophytochromes or phytofluors.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/18326

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) : C12N 9/02, 1/20, 15/00; C07H 21/04

US CL : 435/189, 252.3, 320.1; 536/23.2

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/189, 252.3, 320.1; 536/23.2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	FRANKENBERG, et al. Functional Genomic Analysis of the HY2 Family of Ferredoxin-Dependent Bilin Reductases from Oxygenic Photosynthetic Organisms. The Plant Cell. April 2001, Vol. 13, pages 965-978.	1-11, 33-38



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

07 MARCH 2002

Date of mailing of the international search report

08 May 2002

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

Valerie Bell-Harris for  
TEKCHAND SAIDHA

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet) (July 1998)\*



# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/18926

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  
1-11 and 33-38

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☐

No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

international application No.

PCT/US01/18326

## B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

STN files included medline, caplus, embase, biotechds, scisearch. West for USP, EP, WO. Search terms included - ferredoxin dependent bilin reductase, pebA, PebB, consensus sequence?, phytobilin reductase.

## BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

## THE FOLLOWING CLAIMS ARE GENERIC

Group I, claim(s) 1-11 & 33-38 drawn to consensus sequence (protein) of HY2 family bilin reductase (claims 1-11), and the encoding nucleic acid species in claims 33-38 corresponding to consensus sequence only, host cell and vector.

Group II, claim(s) 12-18 & 33-38, drawn to bilin reductase of 19 reductase species (see Claim 12) and the encoding nucleic acid species in claims 33-38, host cell and vector.

Group III, claim(s) 19-20, drawn to bilin reductase of 23 reductase species (see Claim 19) and the encoding nucleic acid species in claims 33-38, host cell and vector.

Group IV, claim(s) 21-32, drawn to a method of converting biliverdin to phytobilin.

Group V, claim(s) 39, drawn to a method of detecting expression.

Group VI, claim(s) 40-54, drawn to a method of producing photoactive holophytochrome.

Group VII, claim(s) 55-69, drawn to a cell comprising heme oxygenase, an apophytochrome and a ferredoxin-dependent bilin reductase.

Group VIII, claim(s) 70-79, drawn to a recombinant nucleic acid comprising a nucleic acid encoding heme oxidoreductase and a nucleic acid encoding a ferredoxin-dependent bilin reductase.

This application contains claims directed to more than one species of the generic invention. These species are deemed to lack Unity of Invention because they are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for more than one species to be searched, the appropriate additional search fees must be paid. The species are as follows:

As presented above Group II has 19 reductase species and Group III has 23 reductase species; and Groups I, IV, V, VI, VII and VIII have 1 species each and a total of 48 groups.

The inventions listed as Groups I-VIII do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: Group I has a special technical feature of the consensus reductase sequence encoded by a nucleotide sequence, which Groups II-VII do not share; Group II has a specific reductase (species) each encoded by a structurally distinct nucleic acid, which Groups I & III-VIII do not share; Group III has a specific reductase (species) each encoded by a structurally distinct nucleic acid, which Groups I-II & IV-VIII do not share; Groups IV-VI are diverse methods for detecting expression, making phytobilin and for producing photoactive holophytochrome and employ the various reductases and nucleic acids, however, in view of 37 CFR 1.475(b), when claims corresponding to different categories of inventions are present additional methods of use are deemed to lack unity. Groups VII and VIII have a special technical feature of a combination of enzymes or nucleic acids, which Groups VII and VIII do not share among themselves or with Groups I-VI.

Further species presented in Groups II & III are structurally diverse. Thus the various groups and species discussed above show a lack of unity of invention.